

---

# Optimal Algorithms for Decentralized Stochastic Variational Inequalities

---

**Dmitry Kovalev**  
KAUST\*, Saudi Arabia  
dakovalev1@gmail.com

**Aleksandr Beznosikov**  
MIPT†, HSE University and Yandex, Russia  
anbeznosikov@gmail.com

**Abdurakhmon Sadiev**  
MIPT, Russia  
sadiev.aa@phystech.edu

**Michael Persiianov**  
MIPT, Russia  
persiianov.mi@phystech.edu

**Peter Richtárik**  
KAUST, Saudi Arabia  
peter.richtarik@kaust.edu.sa

**Alexander Gasnikov**  
MIPT, HSE University and IITP RAS‡, Russia  
gasnikov@yandex.ru

## Abstract

Variational inequalities are a formalism that includes games, minimization, saddle point, and equilibrium problems as special cases. Methods for variational inequalities are therefore universal approaches for many applied tasks, including machine learning problems. This work concentrates on the decentralized setting, which is increasingly important but not well understood. In particular, we consider decentralized stochastic (sum-type) variational inequalities over fixed and time-varying networks. We present lower complexity bounds for both communication and local iterations and construct optimal algorithms that match these lower bounds. Our algorithms are the best among the available literature not only in the decentralized stochastic case, but also in the decentralized deterministic and non-distributed stochastic cases. Experimental results confirm the effectiveness of the presented algorithms.

## 1 Introduction

Variational inequalities are a broad and flexible class of problems that includes minimization, saddle point, Nash equilibrium, and fixed point problems as special cases; see [24, 6] for an introduction. Over the long history of modern research on variational inequalities spanning at least half a century, the community developed their own methods and theory, differing from the approaches in their sister field, optimization. The ExtraGradient / MirrorProx methods due to [38, 52, 34] have a similar foundational standing in the variational inequalities field that gradient descent occupies in the optimization literature. As in the case of gradient descent, many modifications [31] and variants [66] of these methods were proposed and studied in the variational inequalities literature.

---

\*King Abdullah University of Science and Technology

†Moscow Institute of Physics and Technology

‡Institute for Information Transmission Problems RAS

## 1.1 Applications of variational inequalities

In recent years, there has been a significant increase of research activity in the study of variational inequalities due to new connections to reinforcement learning [58, 32], adversarial training [44], and GANs [28]. In particular, [21, 26, 47, 20, 42, 60] show that even if one considers the classical (in the variational inequalities literature) regime involving monotone and strongly monotone inequalities, it is possible to obtain insights, methods and recommendations useful for the GAN community.

In addition to the above modern applications, and besides their many classical applications in applied mathematics that include economics, equilibrium theory, game theory and optimal control [23], variational inequalities remain popular in supervised learning (with non-separable loss [33]; with non-separable regularizer [3]), unsupervised learning (discriminative clustering [69]; matrix factorization [4]), image denoising [22, 18], robust optimization [9], and non-smooth optimization via smooth reformulations [56, 53].

## 1.2 Processing on the edge

With the proliferation of mobile phones, wearables, digital sensors, smart home appliances, and other devices capable of capturing, storing and processing data, there is an increased appetite to mine the richness contained in these sources for the benefit of humanity. However, at the same time, the traditional centralized approach relying on moving the data into a single proprietary warehouse for processing via suitable machine learning methods is problematic, and a new *modus operandi* is on the rise: processing the data at the source, on the edge, where it was first captured and where it is stored [37, 46, 35], by the client's devices that own the data. There are many reasons for a gradual shift in this direction, including energy efficiency and data privacy.

A key necessary characteristic for any viable algorithmic approach to work in such a massively decentralized regime is the ability to support decentralized processing reflecting the fact that the devices are connected through a network of a potentially complicated topology, possibly varying in time. A central authority may be absent in such a system, and the methods need to rely on communication patterns that correspond to the existing connection links.

## 1.3 Decentralized algorithms for variational inequalities

In this paper we study

*algorithms for solving variational inequalities over decentralized communication networks.*

In this regime, a number of nodes (workers, devices, clients) are connected via a communication network, represented by a graph. Each node can perform computations using its local state and data, and is only allowed to communicate with its neighbors in the graph.

Decentralized algorithms over fixed communication networks find their applications in sensor networks [61], network resource allocation [8], cooperative control [27], distributed spectrum sensing [7], power system control [25] and, of course, in machine learning [63]. Recently, decentralized methods over time-varying networks have gained particular popularity due to their relevance to federated learning [37, 35], where communication failures between devices are a common problem.

Decentralized minimization methods are well studied [49, 36]. In particular, lower bounds and optimal algorithms for such problems are known in the *fixed* [63, 30, 40] and *time-varying* [39, 41] network topology regimes.

*However, in significantly more general and hence potentially much more impactful formalism of variational inequalities, the question of optimal and efficient decentralized methods is still open.*

Motivated by these considerations, our work is devoted to advancing the algorithmic and theoretical foundations of decentralized variational inequalities, in both the fixed and time-varying network regimes.

## 1.4 Our contributions and related work

We now briefly summarize our main contributions.

### (a) Lower bounds

We present the **first lower bounds for the communication and local computation complexities of decentralized variational inequalities in the stochastic (finite sum) case, in both the fixed and time-varying network topology regimes**. See Table 1.

Existing literature contains lower bounds for *non-distributed* finite-sum variational inequalities [29], which we recover as a special case. Existing literature also contains lower bounds for *deterministic* decentralized variational inequalities in the fixed [13] and time-varying [12] regimes. Our bounds covers these results too. See Table 2 (Appendix G).

### (b) Optimal decentralized algorithms

We construct **four new algorithms for stochastic (finite sum) decentralized variational inequalities: two for fixed networks, and two for time-varying networks. Two of these algorithms match our lower bounds, and are therefore optimal in terms of communication and local iteration complexities**. These are the first algorithms for stochastic (finite sum) decentralized variational inequalities over fixed and time-varying networks. See Table 1.

Moreover, our results offer linear communication complexity for deterministic decentralized strongly monotone variational inequalities, which is an improvement upon the sublinear results of [14, 13, 10, 12]. Additionally, our algorithms have better guarantees on local computations than the methods developed by [62]. See Table 2 (Appendix G).

Let us also single out a number of works on decentralized saddle point problems or VIs which are not suitable for comparison with our results: [67, 5] consider non-monotone(minty) problems, [45] does not prove convergence, [43] assumes data homogeneity, and [64] considers a discrete problem.

Table 1: Summary of upper and lower bounds for communication and local computation complexities for finding an  $\varepsilon$ -solution for strongly monotone **stochastic (finite-sum) decentralized** variational inequality (1) over fixed and time-varying networks. Convergence is measured by the distance to the solution.

		Reference	Communication complexity	Local complexity	Weaknesses
Fixed	Upper	Mukherjee and Chakraborty [48] <sup>(1,2)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \frac{D}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(n \chi \frac{L}{\mu} \frac{D}{\mu} \log \frac{1}{\varepsilon}\right)$	weak communication rates weak local computation rates
		Beznosikov et al. [14] <sup>(1,2)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(n \frac{L}{\mu} \log \frac{L+\mu}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip no linear convergence
		Beznosikov et al. [13] <sup>(1,3)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(n \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip no linear convergence
		Rogozin et al. [62] <sup>(1,2,4)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(n \sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	weak local computation rates
		Alg. 1 (this paper)	$\mathcal{O}\left(\max[\sqrt{n}; \sqrt{\chi}] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(6)</sup>	$\mathcal{O}\left(\max[\sqrt{n}; \sqrt{\chi}] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(6)</sup>	
	Alg. 1 + Alg. 3 (this paper)	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(6)</sup>	$\mathcal{O}\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(6)</sup>	multiple gossip	
Lower	Beznosikov et al. [13] <sup>(3)</sup>	$\Omega\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\Omega\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$		
	Thm. 3.2 + Cor. 3.3 (this paper)	$\Omega\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\Omega\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$		
Time-varying	Upper	Beznosikov et al. [10] <sup>(3)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon} + \chi \frac{LD}{\mu^2 \sqrt{\varepsilon}}\right)$ <sup>(5)</sup>	$\mathcal{O}\left(n \chi \frac{L}{\mu} \log \frac{1}{\varepsilon} + n \chi \frac{LD}{\mu^2 \sqrt{\varepsilon}}\right)$	$D$ -homogeneity no linear convergence
		Beznosikov et al. [12] <sup>(1,2)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(n \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip no linear convergence
		Alg. 2 (this paper)	$\mathcal{O}\left(\max[\sqrt{n}; \chi] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(5,6)</sup>	$\mathcal{O}\left(\max[\sqrt{n}; \chi] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(6)</sup>	
	Alg. 2 + Eq. 6 (this paper)	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(5,6)</sup>	$\mathcal{O}\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(6)</sup>	multiple gossip	
	Lower	Beznosikov et al. [12] <sup>(2)</sup>	$\Omega\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\Omega\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	
Thm. 3.4 + Cor. 3.5 (This paper)	$\Omega\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(5)</sup>	$\Omega\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$			

<sup>(1)</sup> for saddle point problems; <sup>(2)</sup> deterministic; <sup>(3)</sup> stochastic, but not finite sum; <sup>(4)</sup> convex-concave (monotone) case (we re-analyzed for strongly monotone case); <sup>(5)</sup>  $B$ -connected graphs [51] are also considered. For simplicity in comparison with other works, we put  $B = 1$ . To get estimates for  $B \neq 1$ , one need to change  $\chi$  to  $B\chi$ ; <sup>(6)</sup> can include additional factors such as  $n \log \frac{1}{\varepsilon}$ ,  $\chi \log \frac{1}{\varepsilon}$ , for full complexities, see details in Section 4.

Notation:  $\mu =$  constant of strong monotonicity of operator  $F$ ,  $L =$  Lipschitz constants of  $L_{m,i}$ ,  $\chi =$  characteristic number of the network (see Assumptions 2.3 and 2.4),  $n =$  size of the local dataset.

### (c) Optimal non-distributed/centralized algorithms

We believe it is notable that despite the generality of our setup and algorithms, **our results, when specialized to handle this simpler case, improve upon the current state-of-the-art results in the non-distributed/centralized setting**. In particular, unlike existing methods, our algorithms support *batching*: while the complexity of the best available algorithms grows with the batch size, our algorithms are not sensitive to this. This property is of crucial importance when working in the large batch mode, which is used in the practice [16, 73, 71]. See Table 3 (Appendix G).

## (d) Experiments

Numerical experiments on bilinear problems and robust regression problems confirm the practical efficiency of our methods, both in the non-distributed stochastic setup and in the decentralized deterministic one.

## 2 Problem Setup and Assumptions

We write  $\langle x, y \rangle := \sum_{i=1}^n x_i y_i$  to denote the standard inner product of vectors  $x, y \in \mathbb{R}^n$ , where  $x_i$  corresponds to the  $i$ -th component of  $x$  in the standard basis in  $\mathbb{R}^n$ . This induces the standard  $\ell_2$ -norm in  $\mathbb{R}^n$  in the following way:  $\|x\| := \sqrt{\langle x, x \rangle}$ . To denote the Kronecker product of two matrices  $\mathbf{A} \in \mathbb{R}^{m \times m}$  and  $\mathbf{B} \in \mathbb{R}^{n \times n}$ , we use  $A \otimes B \in \mathbb{R}^{nm \times nm}$ . The identity matrix of size  $n \times n$  is denoted by  $\mathbf{I}_n$ . We write  $[n] := \{1, 2, \dots, n\}$ .  $\mathbb{N}$  is the set of positive integers.

### 2.1 Variational inequality

We study variational inequalities (VI) of the form

$$\text{Find } z^* \in \mathbb{R}^d \text{ such that } \langle F(z^*), z - z^* \rangle + g(z) - g(z^*) \geq 0, \quad \forall z \in \mathbb{R}^d, \quad (1)$$

where  $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is an operator, and  $g : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$  is a proper lower semicontinuous convex function. We also assume that  $g$  is proximal friendly, i.e. the computation of the operator  $\text{prox}_{\rho g}(z) = \arg \min_{y \in \mathbb{R}^d} \{\rho g(y) + \frac{1}{2} \|y - z\|^2\}$  (with  $\rho > 0$ ) is done for free or costs very low.

To showcase the expressive power of the formalism (1), we now give a few examples of variational inequalities arising in machine learning.

**Example 1 [Convex minimization].** Consider the convex regularized minimization problem:

$$\min_{z \in \mathbb{R}^d} f(z) + g(z), \quad (2)$$

where  $f$  is typically a smooth data-fidelity term, and  $g$  a possibly nonsmooth regularizer. If we define  $F(z) := \nabla f(z)$ , then it can be proved that  $z^* \in \text{dom } g$  is a solution for (1) if and only if  $z^* \in \text{dom } g$  is a solution for (2). So, the regularized optimization problem (2) can be cast as a VI (1).

While minimization problems are widely studied in a separate literature, the next class of problems is much more strongly tied to variational inequalities.

**Example 2 [Convex-concave saddles].** Consider the convex-concave saddle point problem

$$\min_{x \in \mathbb{R}^{d_x}} \max_{y \in \mathbb{R}^{d_y}} f(x, y) + g_1(x) - g_2(y), \quad (3)$$

where  $g_1$  and  $g_2$  can also be interpreted as regularizers. If we let  $F(z) := F(x, y) = [\nabla_x f(x, y), -\nabla_y f(x, y)]$  and  $g(z) = g(x, y) = g_1(x) + g_2(y)$ , then it can be proved that  $z^* \in \text{dom } g$  is a solution for (1) if and only if  $z^* \in \text{dom } g$  is a solution for (3). So, convex-concave saddle point problems (3) can be cast as a VI (1).

Saddle point problems are strongly related to variational inequalities. In particular, lower bounds for the former are also valid for the latter. Moreover, upper bounds for variational inequalities are valid for saddle point problems. However, what is perhaps more important is that these lower and upper bounds match. This is in contrast to minimization, where the lower bounds are weaker.

### 2.2 Decentralized variational inequalities

We consider the decentralized case of problem (1), namely we assume that  $F$  is distributed across  $M$  workers,

$$F(z) := \sum_{m=1}^M F_m(z), \quad (4)$$

while each  $F_m : \mathbb{R}^d \rightarrow \mathbb{R}^d$ ,  $m \in [M]$ , has the finite sum structure

$$F_m(z) := \frac{1}{n} \sum_{i=1}^n F_{m,i}(z). \quad (5)$$

The data describing  $F_m$  being stored on worker  $m$ . For example,  $F_{m,i}$  can correspond to the value of the operator on the  $i$ th data point of the  $m$ -th dataset.

## 2.3 Assumptions

**Assumption 2.1** (Lipschitzness). Each operator  $F_m$  is  $L$ -Lipschitz continuous, i.e. for all  $u, v \in \mathbb{R}^d$  we have  $\|F_m(u) - F_m(v)\| \leq L\|u - v\|$ .

Further, the collection of operators is  $\bar{L}$ -average Lipschitz continuous, i.e., for all  $u, v \in \mathbb{R}^d$  it holds  $\frac{1}{n} \sum_{i=1}^n \|F_{m,i}(u) - F_{m,i}(v)\|^2 \leq \bar{L}^2 \|u - v\|^2$ .

In the context of (2) and (3),  $L$ -Lipschitzness of the operator means that the functions  $f(z)$  and  $f(x, y)$  are  $L$ -smooth.

**Assumption 2.2** (Strong monotonicity). Each operator  $F_m$  is  $\mu$ -strongly monotone, i.e., for all  $u, v \in \mathbb{R}^d$  we have  $\langle F_m(u) - F_m(v); u - v \rangle \geq \mu \|u - v\|^2$ .

In the context of (2) and (3), strong monotonicity of  $F$  means strong convexity of  $f(z)$  and strong convexity-strong concavity of  $f(x, y)$ .

## 2.4 Communication and gossip

Typically, decentralized communication is realized via a *gossip protocol* [68, 15, 50], which is merely matrix-vector multiplication with a gossip matrix  $\mathbf{W}$ , described below, which is different in the fixed and time-varying cases. Let  $\mathcal{L} = \{\mathbf{z} = (z_1, \dots, z_M)^\top \in (\mathbb{R}^d)^M : z_1 = \dots = z_M\}$  be the *consensus space*.

**Assumption 2.3** (Fixed network [63]). For a fixed network, communication can be modeled via an undirected connected graph,  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ , where  $\mathcal{V} = [n]$  are vertices (workers) and  $\mathcal{E} = \{(i, j) \mid i, j \in \mathcal{V}\}$  are edges. Note that  $(i, j) \in \mathcal{E}$  if and only if there exists a communication link between agents  $i$  and  $j$ . The gossip matrix  $\mathbf{W}$  satisfies the following three assumptions: 1)  $\mathbf{W}$  is symmetric positive semi-definite; 2)  $\ker \mathbf{W} \supset \mathcal{L}$ ; 3)  $\mathbf{W}$  is supported on the vertices and edges of the network only:  $w_{i,j} \neq 0$  if and only if  $i = j$  or  $(i, j) \in \mathcal{E}$ .

To characterize the matrix  $\mathbf{W}$ , which captures the properties of the network, we denote  $\lambda_{\max}(\mathbf{W}) = 1$  as the maximum eigenvalue of  $\mathbf{W}$ ,  $\lambda_{\min}^+(\mathbf{W})$  as the minimum positive eigenvalue of  $\mathbf{W}$ , and the characteristic number  $\chi = \lambda_{\max}(\mathbf{W}) / \lambda_{\min}^+(\mathbf{W}) = 1 / \lambda_{\min}^+(\mathbf{W})$ .

**Assumption 2.4** (Time-varying network [51]). For a time-varying network, at any moment  $t$ , communication network can be modeled as a directed  $B$ -connected graph,  $\mathcal{G}(t) = (\mathcal{V}, \mathcal{E}(t))$ , where  $\mathcal{E}(t) = \{(i, j) \mid i, j \in \mathcal{V}\}$  are directed edges.  $B$ -connectedness means that for any time  $t$ , the graph  $\mathcal{G}_B(t)$  with the set of edges  $\bigcup_{\tau=t}^{t+B-1} \mathcal{E}(\tau)$  is connected. To describe the gossip protocol for time-varying case, we define the multi-consensus gossip matrix

$$\mathbf{W}_T(t) = \mathbf{I}_M - \prod_{\tau=t}^{t+T-1} \mathbf{W}(\tau). \quad (6)$$

One can also observe that multiplication with the matrix  $\mathbf{W}_T$  requires to perform multiplication with  $T$  gossip matrices  $\mathbf{W}(t), \dots, \mathbf{W}(t+T-1)$ , i.e., it requires  $T$  decentralized communications. We further assume that the gossip matrices  $\mathbf{W}(t)$  (for  $\mathcal{G}(t)$ ) and  $\mathbf{W}_B(t)$  satisfy: 1)  $\mathbf{W}(t)$  is supported on the nodes and edges of the network:  $w_{i,j}(t) \neq 0$  if and only if  $i = j$  or  $(i, j) \in \mathcal{E}(t)$ ; 2)  $\ker \mathbf{W}(t) \supset \mathcal{L}$ ; 3)  $\text{range} \mathbf{W}(t) \subset \{\mathbf{z} \in (\mathbb{R}^d)^M : \sum_{m=1}^M z_m = 0\}$ ; 4) there exists a characteristic number  $\chi \geq 1$  such that  $\|\mathbf{W}_B(t)z - z\|^2 \leq (1 - \chi^{-1})\|z\|^2$  for all  $z \in \text{range} \mathbf{W}_B(t)$ .

## 3 Lower Bounds

Our lower bounds apply to a specific class of algorithms which are, loosely speaking, allowed to communicate with neighbors, and compute any local first-order information. We now give a formal definition.

**Definition 3.1** (Oracle). Each agent  $m$  has its own local memory  $\mathcal{M}_m$  with initialization  $\mathcal{M}_m = \{0\}$ .  $\mathcal{M}_m$  is updated as follows. At each iteration, the algorithm either performs local computations or communicates.

• **Local computation:** At each local iteration, device  $m$  can sample uniformly and independently batch  $S_m$  of any size  $b$  from  $\{F_{m,i}\}$  and adds to its  $\mathcal{M}_m$  a finite number of points  $z$ , satisfying

$$z \in \text{span} \left\{ z', \sum_{i_m \in S_m} F_{m,i_m}(z''), \text{prox}_{\rho g} \left( \text{span} \left\{ z''', \sum_{i_m \in S_m} F_{m,i_m}(z''') \right\} \right) \right\} \quad (7)$$

for  $z', z'', z''' \in \mathcal{M}_m$  and  $\rho > 0$ . Such a call needs  $b$  local computations to collect the batch. Batch of size  $n$  represents  $F_m$ ;

• **Communication:** Upon communication rounds among the neighboring nodes, and at communication time  $t$ ,  $\mathcal{M}_m$  is updated according to

$$\mathcal{M}_m := \text{span} \left\{ \bigcup_{(i,m) \in \mathcal{E}(t)} \mathcal{M}_i \right\}. \quad (8)$$

• **Output:** The final global output is calculated as  $\hat{z} \in \text{span} \left\{ \bigcup_{m=1}^M \mathcal{M}_m \right\}$ .

The structure of the above definition is typical for distributed lower bounds [63] and for stochastic lower bounds [30]. In particular, Definition 3.1 includes all the approaches for working with stochastic problems, such as SGD or variance reduction techniques (SVRG, SARAH). Note that while our algorithm can invoke the deterministic oracle (full  $F_m$ ) in local computations, in the work on lower bounds in the non-distributed case [29], there is no such a possibility. This narrows the class of algorithms for which results of [29] are valid. In particular, they can not do SVRG-type updates.

**Theorem 3.2** (Lower bound - fixed network). *For any  $\bar{L} \geq \mu > 0$  and  $\chi \geq 1$ ,  $n \in \mathbb{N}$  and  $K, N \in \mathbb{N}$ , there exists a decentralized variational inequality (satisfying Assumptions 2.1 and 2.2) on  $\mathbb{R}^d$  (where  $d$  is sufficiently large) with  $z^* \neq 0$  over a fixed network (satisfying Assumption 2.3) with a gossip matrix  $\mathbf{W}$  and characteristic number  $\chi$ , such that for any output  $\hat{z}$  of any procedure (Definition 3.1) with  $K$  communication rounds and  $N$  local computations, it holds that  $\mathbb{E}[\|\hat{z} - z^*\|^2]$  is*

$$\Omega \left( \exp \left( -\frac{80}{1 + \sqrt{\frac{2\bar{L}^2}{\mu^2} + 1}} \cdot \frac{K}{\sqrt{\chi}} \right) R_0^2 \right) \text{ and } \Omega \left( \exp \left( -\frac{16}{n + \sqrt{\frac{2n\bar{L}^2}{\mu^2} + n^2}} \cdot N \right) R_0^2 \right),$$

where  $R_0^2 = \|z^0 - z^*\|^2$  and  $L = \frac{\bar{L}}{\sqrt{n}}$ .

**Corollary 3.3.** *In the setting of Theorem 3.2, the number of communication rounds and local computations required to obtain an  $\varepsilon$ -solution (in expectation) is lower bounded by*

$$\Omega \left( \sqrt{\chi} \left( 1 + \frac{L}{\mu} \right) \cdot \log \left( \frac{R_0^2}{\varepsilon} \right) \right) \text{ and } \Omega \left( \left( n + \sqrt{n} \cdot \frac{\bar{L}}{\mu} \right) \cdot \log \left( \frac{R_0^2}{\varepsilon} \right) \right), \text{ respectively.}$$

**Theorem 3.4** (Lower bound - time varying network). *For any  $\bar{L} \geq \mu > 0$  and  $\chi \geq 3$ ,  $n \in \mathbb{N}$  and  $K, N \in \mathbb{N}$ , there exist a decentralized variational inequality (satisfying Assumptions 2.1 and 2.2) on  $\mathbb{R}^d$  (where  $d$  is sufficiently large) with  $z^* \neq 0$  over a time-varying network (satisfying Assumption 2.4) with a sequence of gossip matrices  $\mathbf{W}(t)$  and characteristic number  $\chi$ , such that for any output  $\hat{z}$  of any procedure (Definition 3.1) with  $K$  communication rounds and  $N$  local computations, it holds that  $\mathbb{E}[\|\hat{z} - z^*\|^2]$  is*

$$\Omega \left( \exp \left( -\frac{64}{\left( 1 + \sqrt{\frac{2\bar{L}^2}{\mu^2} + 1} \right) \cdot B\chi} \cdot \frac{K}{B\chi} \right) R_0^2 \right) \text{ and } \Omega \left( \exp \left( -\frac{16}{n + \sqrt{\frac{2n\bar{L}^2}{\mu^2} + n^2}} \cdot N \right) R_0^2 \right),$$

where  $R_0^2 = \|z^0 - z^*\|^2$  and  $L = \frac{\bar{L}}{\sqrt{n}}$ .

**Corollary 3.5.** *In the setting of Theorem 3.4, the number of communication rounds and local computations required to obtain an  $\varepsilon$ -solution (in expectation) is lower bounded by*

$$\Omega \left( B\chi \left( 1 + \frac{L}{\mu} \right) \cdot \log \left( \frac{R_0^2}{\varepsilon} \right) \right) \text{ and } \Omega \left( \left( n + \sqrt{n} \cdot \frac{\bar{L}}{\mu} \right) \cdot \log \left( \frac{R_0^2}{\varepsilon} \right) \right), \text{ respectively.}$$

See proofs for Theorems 3.2 and 3.4 in Appendix C. The proof uses the idea (example of bad functions) of non-distributed deterministic lower bounds from [72]. This idea is further extended to distributed stochastic VIs.

Note that in the time-varying case, the lower bounds for communication differ by the constant  $B$  from the estimates that were previously encountered in the literature [12]. This is due to the fact that we consider a more general setup with a  $B$ -connected graph (see Assumption 2.4), while the existing literature on lower bounds focuses on the simpler  $B = 1$  case.

## 4 Optimal Algorithms

Since in the decentralized gossip protocol each local worker  $m$  stores its own  $z_m$  vector, we consider the problem

$$\text{Find } \mathbf{z}^* \in (\mathbb{R}^d)^M \text{ such that } \langle \mathbf{F}(\mathbf{z}^*), \mathbf{z} - \mathbf{z}^* \rangle + \mathbf{g}(\mathbf{z}) - \mathbf{g}(\mathbf{z}^*) \geq 0, \forall \mathbf{z} \in (\mathbb{R}^d)^M, \quad (9)$$

where we use new notation:  $\mathbf{z} = (z_1, \dots, z_M)^\top$  and  $\mathbf{z}^* = (z_1^*, \dots, z_M^*)^\top$ . Additionally, here we introduce the lifted operator  $\mathbf{F} : (\mathbb{R}^d)^M \rightarrow (\mathbb{R}^d)^M$  given as  $\mathbf{F}(\mathbf{z}) = (F_1(z_1), \dots, F_M(z_M))^\top$ , and the lifted operator  $\mathbf{g} : (\mathbb{R}^d)^M \rightarrow \mathbb{R} \cup \{+\infty\}$  defined by  $\mathbf{g}(\mathbf{z}) = \frac{1}{M} \sum_{m=1}^M g(z_m)$ .

One can note that (9) is a set of  $M$  unrelated variational inequalities with their own variables. But the original problem (1) + (5) is a sum of variational inequalities with the same variables:  $\sum_{m=1}^M \left[ \langle F(z_m^*), z_m - z_m^* \rangle + \frac{1}{M} g(z_m) - \frac{1}{M} g(z_m^*) \right]$ . To eliminate this issue and move on to problem (1) + (5), it is easy to get the following modification of (9)

$$\text{Find } \mathbf{z}^* \in \mathcal{L} \text{ such that } \langle \mathbf{F}(\mathbf{z}^*), \mathbf{z} - \mathbf{z}^* \rangle + \mathbf{g}(\mathbf{z}) - \mathbf{g}(\mathbf{z}^*) \geq 0, \forall \mathbf{z} \in \mathcal{L}, \quad (10)$$

where  $\mathcal{L}$  is the consensus space. Problem (10) is equivalent to (1) + (5). Due to Assumptions 2.1 and 2.2,  $\mathbf{F}$  is  $L$ -Lipschitz continuous,  $\bar{L}$ -average Lipschitz continuous and  $\mu$ -strongly monotone.

### 4.1 Fixed networks

We present Algorithm 1 for fixed networks. In Appendix B we give a discussion and intuition. In particular, we give the deterministic variant as well as the non-distributed version of Algorithm 1. The next result gives the iteration complexity of Algorithm 1.

**Theorem 4.1** (Upper bound - fixed network). *Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a fixed graph  $\mathcal{G}$  (Assumption 2.3) with a gossip matrix  $\mathbf{W}$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 1 with tuning of  $\eta, \theta, \alpha, \beta, \gamma$  as described in Appendix D. Then, given  $\varepsilon > 0$ , the number of iterations for  $\mathbb{E}[\|\mathbf{z}^k - \mathbf{z}^*\|^2] \leq \varepsilon$  is*

$$\mathcal{O} \left( \left[ \frac{1}{p} + \chi + \frac{1}{\sqrt{pb}} \frac{\bar{L}}{\mu} + \sqrt{\chi} \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right).$$

See the proof in Appendix D. Let us discuss the results of Theorem. First of all, we are interested in how to obtain the complexity of communications and local computations from iterative complexity. At each iteration we require (in average)  $\mathcal{O}(b + pn)$  local computations, because we need to store batch  $b$  twice and with probability  $p$  we update the point  $\mathbf{w}^{k+1}$  by  $\mathbf{z}^k$ , this requires calculating the full  $\mathbf{F}$  in the next iteration. Then, as the optimal  $p$ , one can choose  $p \sim b/n$ . Then with such choice of  $p$  we have the following local and communication complexities

$$\mathcal{O} \left( \left[ n + b\chi + \sqrt{n} \frac{\bar{L}}{\mu} + b\sqrt{\chi} \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right) \quad \text{and} \quad \mathcal{O} \left( \left[ \frac{n}{b} + \chi + \frac{\sqrt{n}}{b} \frac{\bar{L}}{\mu} + \sqrt{\chi} \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right),$$

respectively (since at each iteration Algorithm 1 performs  $\mathcal{O}(1)$  communications).

Hence, with  $b = 1$  we have the complexities the same as in Table 1. Depending on  $\max\{\sqrt{n}; \sqrt{\chi}\}$ , we have the optimality of either local communications or decentralized communications. One can

---

#### Algorithm 1

---

- 1: **Parameters:** Stepsizes  $\eta, \theta > 0$ , momentums  $\alpha, \beta, \gamma$ , batchsize  $b \in \{1, \dots, n\}$ , probability  $p \in (0, 1)$
- 2: **Initialization:** Choose  $\mathbf{z}^0 = \mathbf{w}^0 \in (\text{dom } g)^M$ ,  $\mathbf{y}^0 \in \mathcal{L}^\perp$ . Put  $\mathbf{z}^{-1} = \mathbf{z}^0$ ,  $\mathbf{w}^{-1} = \mathbf{w}^0$ ,  $\mathbf{y}^{-1} = \mathbf{y}^0$
- 3: **for**  $k = 0, 1, 2 \dots$  **do**
- 4:   Sample  $j_{m,1}^k, \dots, j_{m,b}^k$  independently from  $[n]$
- 5:    $S^k = \{j_{m,1}^k, \dots, j_{m,b}^k\}$
- 6:   Sample  $j_{m,1}^{k+1/2}, \dots, j_{m,b}^{k+1/2}$  independently from  $[n]$
- 7:    $S^{k+1/2} = \{j_{m,1}^{k+1/2}, \dots, j_{m,b}^{k+1/2}\}$
- 8:    $\delta^k = \frac{1}{b} \sum_{j \in S^k} \left( \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1}) \right. \\ \left. + \alpha[\mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1})] \right) + \mathbf{F}(\mathbf{w}^{k-1})$
- 9:    $\Delta^k = \delta^k - (\mathbf{y}^k + \alpha(\mathbf{y}^k - \mathbf{y}^{k-1}))$
- 10:    $\mathbf{z}^{k+1} = \text{prox}_{\eta \mathbf{g}}(\mathbf{z}^k + \gamma(\mathbf{w}^k - \mathbf{z}^k) - \eta \Delta^k)$
- 11:    $\Delta^{k+1/2} = \frac{1}{b} \sum_{j \in S^{k+1/2}} \left( \mathbf{F}_j(\mathbf{z}^{k+1}) - \mathbf{F}_j(\mathbf{w}^k) \right) \\ + \mathbf{F}(\mathbf{w}^k)$
- 12:    $\mathbf{y}^{k+1} = \mathbf{y}^k - \theta(\mathbf{W} \otimes \mathbf{I}_d)(\mathbf{z}^{k+1} - \beta(\Delta^{k+1/2} - \mathbf{y}^k))$
- 13:    $\mathbf{w}^{k+1} = \begin{cases} \mathbf{z}^k, & \text{with probability } p \\ \mathbf{w}^k, & \text{with probability } 1 - p \end{cases}$
- 14: **end for**

---


$$* \mathbf{F}_j(\mathbf{z}) = (F_{1,j_{1,l}}(z_1), \dots, F_{M,j_{M,l}}(z_M))^T, l \in \{1, \dots, b\}$$


---

note that it is enough to take  $b \geq \bar{L}\sqrt{n}/L$  and guarantee the optimal communication complexity (see Corollary 3.3), but we have non-optimality in local iterations.

To make the algorithm optimal both in terms of communications and local computations, we need to slightly modify it. One can make it using Chebyshev acceleration (see Algorithm 3 in Appendix A). Following [63], we can construct a polynomial  $P$  such that 1)  $P(\mathbf{W})$  is a gossip matrix, 2) multiplication by  $P(\mathbf{W}) \otimes \mathbf{I}_d$  requires  $\sqrt{\chi(\mathbf{W})}$  multiplications by  $\mathbf{W}$  (i.e.  $\sqrt{\chi(\mathbf{W})}$  communication rounds) 3)  $\chi(P(\mathbf{W})) \leq 4$ . Then we can modify Algorithm 1 by replacing  $\mathbf{W}$  by  $P(\mathbf{W})$  and get

**Theorem 4.2** (Upper bound - fixed network). *Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a fixed connected graph  $\mathcal{G}$  (Assumption 2.3) with a gossip matrix  $\mathbf{W}$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 1 with Chebyshev polynomial  $P(\mathbf{W})$  as a gossip matrix and with tuning of  $\eta, \theta, \alpha, \beta, \gamma$  as described in Appendix D. Then, given  $\varepsilon > 0$ , the number of iterations for  $\mathbb{E}[\|\mathbf{z}^k - \mathbf{z}^*\|^2] \leq \varepsilon$  is*

$$\mathcal{O}\left(\left[\frac{1}{p} + \frac{1}{\sqrt{pb}}\frac{\bar{L}}{\mu} + \frac{L}{\mu}\right] \log \frac{1}{\varepsilon}\right).$$

In this case, the communication complexity of one iteration is  $\chi$ , and the local complexity (in average) is still  $\mathcal{O}(b + pn)$ . Then with  $p = b/n$  we get the following local and communication complexities

$$\mathcal{O}\left(\left[n + \sqrt{n}\frac{\bar{L}}{\mu} + b\frac{L}{\mu}\right] \log \frac{1}{\varepsilon}\right) \quad \text{and} \quad \mathcal{O}\left(\left[\sqrt{\chi}\frac{n}{b} + \sqrt{\chi}\frac{\sqrt{n}}{b}\frac{\bar{L}}{\mu} + \sqrt{\chi}\frac{L}{\mu}\right] \log \frac{1}{\varepsilon}\right),$$

respectively. To get optimal results from Table 1 we just need to take  $b = \bar{L}\sqrt{n}/L$ .

In contrast to algorithms of [14, 13] (the closest papers in theoretical convergence), our Algorithm 1 needs multi-consensus/Chebyshev acceleration for both optimal rates, but can work without these additional procedures. Algorithms [14, 13] requires  $\mathcal{O}(\sqrt{\chi} \log \varepsilon^{-1})$  iterations for Chebyshev acceleration, which makes the algorithms less practical.

## 4.2 Time-varying networks

We present Algorithm 2 for time-varying networks. It needs to compute  $\mathbf{W}_T$  using (6), it requires  $T$  communications. In Appendix B we give a discussion and intuition of this algorithm. The next result gives the iteration complexity of Algorithm 2.

**Theorem 4.3** (Upper bound - time varying network). *Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a sequence of time-varying graphs  $\mathcal{G}(k)$  (Assumption 2.4) with gossip matrices  $\mathbf{W}(k)$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 2 with  $T \geq B$  and tuning of parameters as described in Appendix E. Let the choice of  $T$  guarantees contraction property (Assumption 2.4 point 4) with  $\chi(T)$ . Then, given  $\varepsilon > 0$ , the number of iterations for  $\mathbb{E}[\|\mathbf{z}^k - \mathbf{z}^*\|^2] \leq \varepsilon$  is*

$$\tilde{\mathcal{O}}\left(\chi^2(T) + \frac{1}{p} + \chi(T)\frac{L}{\mu} + \frac{1}{\sqrt{bp}}\frac{\bar{L}}{\mu}\right).$$

---

### Algorithm 2

---

- 1: **Parameters:** Stepsizes  $\eta_z, \eta_y, \eta_x, \theta > 0$ , momentums  $\alpha, \gamma, \omega, \tau$ , parameters  $\nu, \beta$ , batchsize  $b \in \{1, \dots, n\}$ , probability  $p \in (0, 1)$
  - 2: **Initialization:** Choose  $\mathbf{z}^0 = \mathbf{w}^0 \in (\text{dom } g)^M$ ,  $\mathbf{y}^0 \in (\mathbb{R}^d)^M$ ,  $\mathbf{x}^0 \in \mathcal{L}^\perp$ . Put  $\mathbf{z}^{-1} = \mathbf{z}^0$ ,  $\mathbf{w}^{-1} = \mathbf{w}^0$ ,  $\mathbf{y}_f^{-1} = \mathbf{y}^0$ ,  $\mathbf{x}_f^{-1} = \mathbf{x}^0$ ,  $m_0 = \mathbf{0}^{dM}$
  - 3: **for**  $k = 0, 1, 2, \dots$  **do**
  - 4:   Sample  $j_{m,1}^k, \dots, j_{m,b}^k$  independently from  $[n]$
  - 5:    $S^k = \{j_{m,1}^k, \dots, j_{m,b}^k\}$
  - 6:   Sample  $j_{m,1}^{k+1/2}, \dots, j_{m,b}^{k+1/2}$  independently from  $[n]$
  - 7:    $S^{k+1/2} = \{j_{m,1}^{k+1/2}, \dots, j_{m,b}^{k+1/2}\}$
  - 8:    $\delta^k = \frac{1}{b} \sum_{j \in S^k} \left( \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1}) \right. \\ \left. + \alpha[\mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1})] \right) + \mathbf{F}(\mathbf{w}^{k-1})$
  - 9:    $\Delta_z^k = \delta^k - \nu \mathbf{z}^k - \mathbf{y}^k - \alpha(\mathbf{y}^k - \mathbf{y}^{k-1})$
  - 10:    $\mathbf{z}^{k+1} = \text{prox}_{\eta_z \mathbf{g}}(\mathbf{z}^k + \omega(\mathbf{w}^k - \mathbf{z}^k) - \eta_z \Delta_z^k)$
  - 11:    $\mathbf{y}_c^k = \tau \mathbf{y}^k + (1 - \tau) \mathbf{y}_f^k$
  - 12:    $\mathbf{x}_c^k = \tau \mathbf{x}^k + (1 - \tau) \mathbf{x}_f^k$
  - 13:    $\Delta_y^k = \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \mathbf{z}^{k+1} + \gamma(\mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k)$
  - 14:    $\delta^{k+1/2} = \frac{1}{b} \sum_{j \in S^{k+1/2}} \left( \mathbf{F}_j(\mathbf{z}^{k+1}) - \mathbf{F}_j(\mathbf{w}^k) \right) \\ + \mathbf{F}(\mathbf{w}^k)$
  - 15:    $\Delta_x^k = \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \beta(\mathbf{x}^k + \delta^{k+1/2})$
  - 16:    $\mathbf{y}^{k+1} = \mathbf{y}^k - \eta_y \Delta_y^k$
  - 17:    $\mathbf{x}^{k+1} = \mathbf{x}^k - (\mathbf{W}_T(Tk) \otimes \mathbf{I}_d)(\eta_x \Delta_x^k + m^k)$
  - 18:    $m^{k+1} = \eta_x \Delta_x^k + m^k \\ - (\mathbf{W}_T(Tk) \otimes \mathbf{I}_d)(\eta_x \Delta_x^k + m^k)$
  - 19:    $\mathbf{y}_f^{k+1} = \mathbf{y}_c^k + \tau(\mathbf{y}^{k+1} - \mathbf{y}^k)$
  - 20:    $\mathbf{x}_f^{k+1} = \mathbf{x}_c^k - \theta(\mathbf{W}_T(Tk) \otimes \mathbf{I}_d)(\mathbf{y}^k + \mathbf{x}^k)$
  - 21:    $\mathbf{w}^{k+1} = \begin{cases} \mathbf{z}^k, & \text{with probability } p \\ \mathbf{w}^k, & \text{with probability } 1 - p \end{cases}$
  - 22: **end for**
-



Note that an important detail of the method is that  $T \geq B$ . This limitation is due to the fact that network is  $B$ -connected. In particular, for  $B > 1$  it can happen that in some communications we use empty graphs. Therefore, the requirement  $T \geq B$  is natural to guarantee the contraction property (Assumption 2.4, point 4). This means that if  $B > 1$  we have to use multi-consensus (6).

But with  $B = T = 1$ , we can avoid multi-consensus, let us this case first. The same way as in fixed graph case we choose  $p = b/n$ . Then we get the estimates on communications and local calls

$$\mathcal{O} \left( \left[ \chi^2 + \frac{n}{b} + \chi \frac{L}{\mu} + \frac{\sqrt{n}}{b} \frac{\bar{L}}{\mu} \right] \log \frac{1}{\varepsilon} \right).$$

If we put  $b = 1$ , we have the same estimates as in Table 1.

Now we consider general case with any  $B$ . We use a multi-gossip step and take  $T > 1$ . In particular, let us choose  $T = B \cdot \lceil \chi \ln 2 \rceil$ . Than using (6) and point 4 of Assumption 2.4, we can guarantee that  $\|\mathbf{W}_T(t)z - z\|^2 \leq \frac{1}{2}\|z\|^2$ . Therefore,  $\chi(T) = 2$ , but we need  $T$  communications per iteration. With  $p = b/n$  and  $b = \bar{L}\sqrt{n}/L$ , the iteration complexity from Theorem 4.3 can be rewritten as follows

$$\mathcal{O} \left( \left[ 1 + \frac{\sqrt{n}L}{L} + \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right).$$

Using that per iteration we make  $\mathcal{O}(B \cdot \lceil \chi \ln 2 \rceil)$  communications and  $\mathcal{O}(\bar{L}\sqrt{n}/L)$  local computations, we get

$$\mathcal{O} \left( \left[ B\chi + B\chi \frac{\sqrt{n}L}{L} + B\chi \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right) \text{ comm-n-s and } \mathcal{O} \left( \left[ n + \frac{\bar{L}\sqrt{n}}{L} + \sqrt{n} \frac{\bar{L}}{\mu} \right] \log \frac{1}{\varepsilon} \right) \text{ local calls.}$$

These results are reflected in Table 1.

## 5 Experiments

We now perform several experiments with the goal of corroborating our theoretical results. Note though that we are the first who consider the decentralized stochastic (finite-sum) setting for VIs, and hence there are no competing methods. Therefore, we compare the non-distributed finite sum setting and decentralized deterministic setting separately.

### 5.1 Variance reduction

In this section, we compare the main methods for solving strongly monotone non-distributed stochastic (finite-sum) variational inequalities with non-distributed version of our Algorithm 1.

**Problem.** We first consider bilinear problem:


$$\min_{x \in \Delta^d} \max_{y \in \Delta^d} \frac{1}{n} \sum_{i=1}^n x^\top \mathbf{A}_i y, \quad (11)$$

where  $\Delta^d$  is the unit simplex in  $\mathbb{R}^d$ . We use the same experimental setup as in [1], in particular we consider policeman and burglar matrix from [54] and two test matrices from [55].

**Setting.** For comparison, we took methods from Table 3. In particular, we chose EG-Alc-Alg1 and EG-Alc-Alg2 from [1], EG-Car from [17]. The parameters of all methods are selected in two ways: 1) as described in the theory of the corresponding papers, and 2) tuned for the best convergence. We run all methods with different batch sizes. The comparison criterion is the number of epochs (one full gradient = epoch).

**Results.** The plots from Figure 1 show that in the case of a theoretical choice of parameters, our Algorithm 1 is ahead of other methods for any batch size (including  $b = 1$ ). In the case of tuning parameters, the specialized method from [17] is better than our algorithm. See more experiments with other matrices in Appendix F.1.

Figure 1: Comparison epoch complexities of Algorithm 1, EG-Alc-Alg1, EG-Alc-Alg2 and EG-Car on (11) with matrix from [54]. Dashed lines give convergence with theoretical parameters, solid lines – with tuned parameters.



## 5.2 Decentralized methods

In this section, we compare the state-of-the-art methods for solving strongly monotone decentralized variational inequalities over fixed and time-varying networks with our Algorithms 1 and 2.

**Problem.** We now consider robust linear regression:

$$\min_w \max_{\|r\| \leq \epsilon} \frac{1}{N} \sum_{i=1}^N (w^\top (x_i + r_i) - y_i)^2 + \frac{\lambda}{2} \|w\|^2 - \frac{\beta}{2} \|r\|^2, \quad (12)$$

where  $w$  are model weights,  $\{(x_i, y_i)\}_{i=1}^N$  are pairs of the training data,  $r_i$  are noise vectors, and  $\lambda$  and  $\beta$  are regularization parameters. The noises  $r_i$  resist training the model, thereby inducing more robustness and stability.


**Setting.** For comparison, we took methods from Table 1 for decentralized problems over fixed and time-varying networks. In particular, we choose EGD-GT from [48], EGD-Con from [13, 12] and Sliding from [14]. Note that only EGD-Con has a theory for fixed and time-varying networks, despite this, we use all methods in both cases.

For a fair comparison, we consider the deterministic setup, i.e., each worker can compute full gradients. We take datasets from LIBSVM [19] and divided unevenly across  $M = 25$  workers. For communication networks we chose the star, the ring and the grid topologies. For time-varying networks, the topologies remain the same, but the locations of the vertices in them change randomly. All methods are tuned for the best convergence. The comparison criterion is the number of communication rounds.

### 5.2.1 Fixed networks

**Results.** The plots from Figure 2 show that our Algorithm 1 is ahead of other methods. Among other things, it is ahead of Sliding from [14], which has a fast theoretical communication complexity. However, this happens when the dataset is relatively homogeneous and uniformly divided across the devices. In our setting, this is not the case. See more experiments with other datasets in Appendix F.2.1.


Figure 2: Comparison communication complexities of Algorithm 1, EGD-GT, EGD-Con and Sliding on (12) over fixed networks.



### 5.2.2 Time-varying networks

**Results.** The plots from Figure 3 show that our Algorithm 2 is ahead of other methods in the case of the grid network. In the case of the ring topology, EGD-Con shows the best results. Indeed, such an algorithm in fact implements centralized communications via a decentralized protocol and this approach is not always the fastest, but reliable. See more experiments with other datasets in Appendix F.2.2.

Figure 3: Comparison communication complexities of Algorithm 2, EGD-GT, EGD-Con and Sliding on (12) over time-varying networks.



## Acknowledgments

This work was supported by a grant for research centers in the field of artificial intelligence, provided by the Analytical Center for the Government of the Russian Federation in accordance with the subsidy agreement (agreement identifier 000000D730321P5Q0002) and the agreement with the Moscow Institute of Physics and Technology dated November 1, 2021 No. 70-2021-00138.

## References

- [1] Ahmet Alacaoglu and Yura Malitsky. Stochastic variance reduction for variational inequality methods. *arXiv preprint arXiv:2102.08352*, 2021.

- [2] Ahmet Alacaoglu, Yura Malitsky, and Volkan Cevher. Forward-reflected-backward method with variance reduction. *Computational Optimization and Applications*, 80, 11 2021.
- [3] Francis Bach, Rodolphe Jenatton, Julien Mairal, and Guillaume Obozinski. Optimization with sparsity-inducing penalties. *arXiv preprint arXiv:1108.0775*, 2011.
- [4] Francis Bach, Julien Mairal, and Jean Ponce. Convex sparse matrix factorizations. *arXiv preprint arXiv:0812.1869*, 2008.
- [5] Babak Barzandeh, Tianjian Huang, and George Michailidis. A decentralized adaptive momentum method for solving a class of min-max optimization problems. *Signal Processing*, 189:108245, 2021.
- [6] Heinz Bauschke and Patrick Combettes. *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. 01 2017.
- [7] Juan Andrés Bazerque and Georgios B Giannakis. Distributed spectrum sensing for cognitive radio networks by exploiting sparsity. *IEEE Transactions on Signal Processing*, 58(3):1847–1862, 2009.
- [8] Amir Beck, Angelia Nedić, Asuman Ozdaglar, and Marc Teboulle. An  $o(1/k)$  gradient method for network resource allocation problems. *IEEE Transactions on Control of Network Systems*, 1(1):64–73, 2014.
- [9] Aharon Ben-Tal, Laurent El Ghaoui, and Arkadi Nemirovski. *Robust Optimization*. Princeton University Press, 2009.
- [10] Aleksandr Beznosikov, Pavel Dvurechensky, Anastasia Koloskova, Valentin Samokhin, Sebastian U Stich, and Alexander Gasnikov. Decentralized local stochastic extra-gradient for variational inequalities. *arXiv preprint arXiv:2106.08315*, 2021.
- [11] Aleksandr Beznosikov, Eduard Gorbunov, Hugo Berard, and Nicolas Loizou. Stochastic gradient descent-ascent: Unified theory and new efficient methods. *arXiv preprint arXiv:2202.07262*, 2022.
- [12] Aleksandr Beznosikov, Alexander Rogozin, Dmitry Kovalev, and Alexander Gasnikov. Near-optimal decentralized algorithms for saddle point problems over time-varying networks. *Lecture Notes in Computer Science*, page 246–257, 2021.
- [13] Aleksandr Beznosikov, Valentin Samokhin, and Alexander Gasnikov. Distributed saddle-point problems: Lower bounds, optimal algorithms and federated GANs. *arXiv preprint arXiv:2010.13112*, 2020.
- [14] Aleksandr Beznosikov, Gesualdo Scutari, Alexander Rogozin, and Alexander Gasnikov. Distributed saddle-point problems under data similarity. *Advances in Neural Information Processing Systems*, 34, 2021.
- [15] Stephen Boyd, Arpita Ghosh, Balaji Prabhakar, and Devavrat Shah. Randomized gossip algorithms. *IEEE transactions on information theory*, 52(6):2508–2530, 2006.
- [16] Andrew Brock, Jeff Donahue, and Karen Simonyan. Large scale gan training for high fidelity natural image synthesis. *arXiv preprint arXiv:1809.11096*, 2018.
- [17] Yair Carmon, Yujia Jin, Aaron Sidford, and Kevin Tian. Variance reduction for matrix games. *arXiv preprint arXiv:1907.02056*, 2019.
- [18] Antonin Chambolle and Thomas Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of mathematical imaging and vision*, 40(1):120–145, 2011.
- [19] Chih-Chung Chang and Chih-Jen Lin. Libsvm: a library for support vector machines. *ACM transactions on intelligent systems and technology (TIST)*, 2(3):1–27, 2011.

- [20] Tatjana Chavdarova, Gauthier Gidel, François Fleuret, and Simon Lacoste-Julien. Reducing noise in gan training with variance reduced extragradient. *arXiv preprint arXiv:1904.08598*, 2019.
- [21] Constantinos Daskalakis, Andrew Ilyas, Vasilis Syrgkanis, and Haoyang Zeng. Training gans with optimism. *arXiv preprint arXiv:1711.00141*, 2017.
- [22] Ernie Esser, Xiaoqun Zhang, and Tony F Chan. A general framework for a class of first order primal-dual algorithms for convex optimization in imaging science. *SIAM Journal on Imaging Sciences*, 3(4):1015–1046, 2010.
- [23] F. Facchinei and J.S. Pang. *Finite-Dimensional Variational Inequalities and Complementarity Problems*. Springer Series in Operations Research and Financial Engineering. Springer New York, 2007.
- [24] Francisco Facchinei and Jong-Shi Pang. *Finite-Dimensional Variational Inequalities and Complementarity Problems*. Springer Series in Operations Research. Springer, 2003.
- [25] Lingwen Gan, Ufuk Topcu, and Steven H Low. Optimal decentralized protocol for electric vehicle charging. *IEEE Transactions on Power Systems*, 28(2):940–951, 2012.
- [26] Gauthier Gidel, Hugo Berard, Gaëtan Vignoud, Pascal Vincent, and Simon Lacoste-Julien. A variational inequality perspective on generative adversarial networks. *arXiv preprint arXiv:1802.10551*, 2018.
- [27] Pontus Giselsson, Minh Dang Doan, Tamás Keviczky, Bart De Schutter, and Anders Rantzer. Accelerated gradient methods and dual decomposition in distributed model predictive control. *Automatica*, 49(3):829–833, 2013.
- [28] Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. In *Advances in Neural Information Processing Systems (NIPS)*, 2014.
- [29] Yuze Han, Guangzeng Xie, and Zhihua Zhang. Lower complexity bounds of finite-sum optimization problems: The results and construction. *arXiv preprint arXiv:2103.08280*, 2021.
- [30] Hadrien Hendrikx, Francis Bach, and Laurent Massoulié. An optimal algorithm for decentralized finite sum optimization. *arXiv preprint arXiv:2005.10675*, 2020.
- [31] Yu-Guan Hsieh, Franck Iutzeler, Jérôme Malick, and Panayotis Mertikopoulos. On the convergence of single-call stochastic extra-gradient methods. *arXiv preprint arXiv:1908.08465*, 2019.
- [32] Yujia Jin and Aaron Sidford. Efficiently solving MDPs with stochastic mirror descent. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, volume 119, pages 4890–4900. PMLR, 2020.
- [33] Thorsten Joachims. A support vector method for multivariate performance measures. pages 377–384, 01 2005.
- [34] Anatoli Juditsky, Arkadii S. Nemirovskii, and Claire Tauvel. Solving variational inequalities with stochastic mirror-prox algorithm, 2008.
- [35] Peter Kairouz, H Brendan McMahan, Brendan Avent, Aurélien Bellet, Mehdi Bennis, Arjun Nitin Bhagoji, Kallista Bonawitz, Zachary Charles, Graham Cormode, Rachel Cummings, et al. Advances and open problems in federated learning. *arXiv preprint arXiv:1912.04977*, 2019.
- [36] Anastasia Koloskova, Nicolas Loizou, Sadra Boreiri, Martin Jaggi, and Sebastian U Stich. A unified theory of decentralized sgd with changing topology and local updates. *arXiv preprint arXiv:2003.10422*, 2020.
- [37] Jakub Konečný, H. Brendan McMahan, Felix Yu, Peter Richtárik, Ananda Theertha Suresh, and Dave Bacon. Federated learning: strategies for improving communication efficiency. In *NIPS Private Multi-Party Machine Learning Workshop*, 2016.

- [38] G. M. Korpelevich. The extragradient method for finding saddle points and other problems. *Matecon*, 12:747–756, 1976.
- [39] Dmitry Kovalev, Elnur Gasanov, Peter Richtárik, and Alexander Gasnikov. Lower bounds and optimal algorithms for smooth and strongly convex decentralized optimization over time-varying networks. *arXiv preprint arXiv:2106.04469*, 2021.
- [40] Dmitry Kovalev, Adil Salim, and Peter Richtárik. Optimal and practical algorithms for smooth and strongly convex decentralized optimization. *Advances in Neural Information Processing Systems*, 33, 2020.
- [41] Huan Li and Zhouchen Lin. Accelerated gradient tracking over time-varying graphs for decentralized optimization. *arXiv preprint arXiv:2104.02596*, 2021.
- [42] Tengyuan Liang and James Stokes. Interaction matters: A note on non-asymptotic local convergence of generative adversarial networks. In Kamalika Chaudhuri and Masashi Sugiyama, editors, *Proceedings of the Twenty-Second International Conference on Artificial Intelligence and Statistics*, volume 89 of *Proceedings of Machine Learning Research*, pages 907–915. PMLR, 16–18 Apr 2019.
- [43] Mingrui Liu, Wei Zhang, Youssef Mroueh, Xiaodong Cui, Jerret Ross, Tianbao Yang, and Payel Das. A decentralized parallel algorithm for training generative adversarial nets. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2020.
- [44] Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. In *International Conference on Learning Representations (ICLR)*, 2018.
- [45] David Mateos-Núñez and Jorge Cortés. Distributed subgradient methods for saddle-point problems. In *2015 54th IEEE Conference on Decision and Control (CDC)*, pages 5462–5467, 2015.
- [46] H Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Agüera y Arcas. Communication-efficient learning of deep networks from decentralized data. In *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2017.
- [47] Panayotis Mertikopoulos, Bruno Lecouat, Houssam Zenati, Chuan-Sheng Foo, Vijay Chandrasekhar, and Georgios Piliouras. Optimistic mirror descent in saddle-point problems: Going the extra (gradient) mile. *arXiv preprint arXiv:1807.02629*, 2018.
- [48] Soham Mukherjee and Mrityunjay Chakraborty. A decentralized algorithm for large scale min-max problems. In *2020 59th IEEE Conference on Decision and Control (CDC)*, pages 2967–2972, 2020.
- [49] Angelia Nedic. Distributed gradient methods for convex machine learning problems in networks: Distributed optimization. *IEEE Signal Processing Magazine*, 37(3):92–101, 2020.
- [50] Angelia Nedić and Asuman Ozdaglar. Distributed subgradient methods for multi-agent optimization. *IEEE Transactions on Automatic Control*, 54(1):48–61, 2009.
- [51] Angelia Nedich, Alex Olshevsky, and Wei Shi. A geometrically convergent method for distributed optimization over time-varying graphs. In *2016 IEEE 55th Conference on Decision and Control (CDC)*, pages 1023–1029. IEEE, 2016.
- [52] Arkadi Nemirovski. Prox-method with rate of convergence  $o(1/t)$  for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15:229–251, 01 2004.
- [53] Arkadi Nemirovski. Prox-method with rate of convergence  $O(1/t)$  for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15(1):229–251, 2004.
- [54] Arkadi Nemirovski. Mini-course on convex programming algorithms, 2013.

- [55] Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on optimization*, 19(4):1574–1609, 2009.
- [56] Yu Nesterov. Smooth minimization of non-smooth functions. *Mathematical programming*, 103(1):127–152, 2005.
- [57] Yurii E Nesterov. A method for solving the convex programming problem with convergence rate  $o(1/k^2)$ . In *Dokl. akad. nauk Sssr*, volume 269, pages 543–547, 1983.
- [58] Shayegan Omidshafiei, Jason Pazis, Christopher Amato, Jonathan P. How, and John Vian. Deep decentralized multi-task multi-agent reinforcement learning under partial observability. In *Proceedings of the 34th International Conference on Machine Learning (ICML)*, volume 70, pages 2681–2690. PMLR, 2017.
- [59] Balamurugan Palaniappan and Francis Bach. Stochastic variance reduction methods for saddle-point problems. In *Advances in Neural Information Processing Systems*, pages 1416–1424, 2016.
- [60] Wei Peng, Yu-Hong Dai, Hui Zhang, and Lizhi Cheng. Training gans with centripetal acceleration. *Optimization Methods and Software*, 35(5):955–973, 2020.
- [61] M. Rabbat and R. Nowak. Distributed optimization in sensor networks. In *Third International Symposium on Information Processing in Sensor Networks, 2004. IPSN 2004*, pages 20–27, 2004.
- [62] Alexander Rogozin, Alexander Beznosikov, Darina Dvinskikh, Dmitry Kovalev, Pavel Dvurechensky, and Alexander Gasnikov. Decentralized distributed optimization for saddle point problems. *arXiv preprint arXiv:2102.07758*, 2021.
- [63] Kevin Scaman, Francis Bach, Sébastien Bubeck, Yin Tat Lee, and Laurent Massoulié. Optimal algorithms for smooth and strongly convex distributed optimization in networks. *arXiv preprint arXiv:1702.08704*, 2017.
- [64] Kunal Srivastava, Angelia Nedić, and Dušan Stipanović. Distributed min-max optimization in networks. In *2011 17th International Conference on Digital Signal Processing (DSP)*, pages 1–8, 2011.
- [65] Vladislav Tominin, Yaroslav Tominin, Ekaterina Borodich, Dmitry Kovalev, Alexander Gasnikov, and Pavel Dvurechensky. On accelerated methods for saddle-point problems with composite structure. *arXiv preprint arXiv:2103.09344*, 2021.
- [66] Paul Tseng. A modified forward-backward splitting method for maximal monotone mappings. *SIAM Journal on Control and Optimization*, 38(2):431–446, 2000.
- [67] Wenhan Xian, Feihu Huang, Yanfu Zhang, and Heng Huang. A faster decentralized algorithm for nonconvex minimax problems. In M. Ranzato, A. Beygelzimer, Y. Dauphin, P.S. Liang, and J. Wortman Vaughan, editors, *Advances in Neural Information Processing Systems*, volume 34, pages 25865–25877. Curran Associates, Inc., 2021.
- [68] Lin Xiao and Stephen Boyd. Fast linear iterations for distributed averaging. *Systems and Control Letters*, 53(1):65–78, 2004.
- [69] Linli Xu, James Neufeld, Bryce Larson, and Dale Schuurmans. Maximum margin clustering. In L. Saul, Y. Weiss, and L. Bottou, editors, *Advances in Neural Information Processing Systems*, volume 17. MIT Press, 2005.
- [70] Junchi Yang, Negar Kiyavash, and Niao He. Global convergence and variance-reduced optimization for a class of nonconvex-nonconcave minimax problems. *arXiv preprint arXiv:2002.09621*, 2020.
- [71] Yang You, Jing Li, Sashank Reddi, Jonathan Hseu, Sanjiv Kumar, Srinadh Bhojanapalli, Xiaodan Song, James Demmel, Kurt Keutzer, and Cho-Jui Hsieh. Large batch optimization for deep learning: Training bert in 76 minutes. *arXiv preprint arXiv:1904.00962*, 2019.

- [72] Junyu Zhang, Mingyi Hong, and Shuzhong Zhang. On lower iteration complexity bounds for the saddle point problems. *arXiv preprint arXiv:1912.07481*, 2019.
- [73] Chen Zhu, Yu Cheng, Zhe Gan, Siqi Sun, Tom Goldstein, and Jingjing Liu. FreeLB: Enhanced adversarial training for natural language understanding. *arXiv preprint arXiv:1909.11764*, 2019.

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Applications of variational inequalities . . . . .	2
1.2	Processing on the edge . . . . .	2
1.3	Decentralized algorithms for variational inequalities . . . . .	2
1.4	Our contributions and related work . . . . .	2
<b>2</b>	<b>Problem Setup and Assumptions</b>	<b>4</b>
2.1	Variational inequality . . . . .	4
2.2	Decentralized variational inequalities . . . . .	4
2.3	Assumptions . . . . .	5
2.4	Communication and gossip . . . . .	5
<b>3</b>	<b>Lower Bounds</b>	<b>5</b>
<b>4</b>	<b>Optimal Algorithms</b>	<b>7</b>
4.1	Fixed networks . . . . .	7
4.2	Time-varying networks . . . . .	8
<b>5</b>	<b>Experiments</b>	<b>9</b>
5.1	Variance reduction . . . . .	9
5.2	Decentralized methods . . . . .	10
5.2.1	Fixed networks . . . . .	10
5.2.2	Time-varying networks . . . . .	10
<b>A</b>	<b>Chebyshev acceleration</b>	<b>17</b>
<b>B</b>	<b>Discussion and intuition of Algorithms 1 and 2</b>	<b>17</b>
<b>C</b>	<b>Proof of Theorems 3.2 and 3.4</b>	<b>19</b>
C.1	Fixed network . . . . .	24
C.2	Time-varying network . . . . .	26
<b>D</b>	<b>Proof of Theorem 4.1</b>	<b>26</b>
<b>E</b>	<b>Proof of Theorem 4.3</b>	<b>36</b>
<b>F</b>	<b>Additional experiments</b>	<b>53</b>
F.1	Variance reduction . . . . .	53
F.2	Decentralized methods . . . . .	53
F.2.1	Fixed networks . . . . .	53
F.2.2	Time-varying networks . . . . .	55
<b>G</b>	<b>Tables</b>	<b>56</b>



# Supplementary Materials

## A Chebyshev acceleration

---

### Algorithm 3 Chebyshev gossip subroutine [63]

---

1: **Input:**  $\mathbf{z}, \mathbf{W}$   
2:  $c_2 = \frac{\chi+1}{\chi-1}, a_0 = 1, a_1 = c_2, c_3 = \frac{2}{\lambda_{\max}(\mathbf{W}) + \lambda_{\min}^+(\mathbf{W})}$ .  
3:  $\mathbf{z}^0 = \mathbf{z}, \mathbf{z}^1 = c_2(\mathbf{I} - c_3\mathbf{W})\mathbf{z}$ .  
4: **for**  $k = 1, \dots, K - 1$  **do**  
5:    $a_{k+1} = 2c_2a_k - a_{k-1}$ .  
6:    $\mathbf{z}^{k+1} = 2c_2(\mathbf{I} - c_3\mathbf{W})\mathbf{z}^k - \mathbf{z}^{k-1}$ .  
7: **end for**  
8: **Output:**  $\mathbf{z}^0 - \frac{\mathbf{z}^K}{a_K}$ .

---

## B Discussion and intuition of Algorithms 1 and 2

The design of Algorithm 1 and Algorithm 2 is based on a combination of two optimization ideas: variance reduction for variational inequalities (Algorithm 4) and optimal algorithms for decentralized distributed optimization on fixed (Algorithm 5) and time-varying (Algorithm 6) networks.

**Variance reduction for variational inequalities.** The variance reduction technique used in the design of Algorithms 1 and 2 is based on the FoRB with variance reduction [1, 2]. However, there are crucial differences with the FoRB algorithm that we now describe. Consider the variance reduced “gradient” estimator of FoRB in the case of batch size being equal to 1 ( $b = 1$ ). It can be written as follows:

$$\delta^k = F_j(z^k) - F_j(w^{k-1}) + F(w^k), \quad (13)$$

where  $j$  is sampled from  $\{1, \dots, n\}$  uniformly at random. This step in some sense combines the variance reduction technique and so-called “optimistic” step, which leads to a bad convergence rate  $\mathcal{O}\left(n + \sqrt{bn} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$  in the single node setting (see Table 3). Our Algorithm 4 (basic for Algorithm 1 and Algorithm 2) is different. They use the following “gradient” estimator (see line 6 of Algorithm 4):

$$\delta^k = \underbrace{F_j(z^k) - F_j(w^{k-1}) + F(w^{k-1})}_{\text{variance reduction}} + \alpha \underbrace{\left(F_j(z^k) - F_j(z^{k-1})\right)}_{\text{“optimistic” step}}. \quad (14)$$

One can observe that our gradient estimator in some sense separates the variance reduction step and the “optimistic” step. This leads to the better convergence rate  $\mathcal{O}\left(n + \sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$  in the single node setting (see Table 3) and allows to develop the optimal algorithms in the decentralized distributed setting.

**Decentralized distributed optimization.** To provide the optimal algorithms for decentralized stochastic variational inequalities, we use the optimal algorithms for solving decentralized distributed minimization problems for fixed networks [40] and for time-varying networks [39]. These algorithms use Nesterov acceleration [57] which we replace with our “gradient” estimator (14). Another important difference is that we extend our results to the composite case, i.e.,  $g(z) \neq 0$  in the main problem (1) with the help of the proximal operator  $\text{prox}_g(\cdot)$ . This is one of the important contributions of our paper. To the best of our knowledge, optimal algorithms in the decentralized distributed setting exist neither for minimization problems nor for the variational inequalities.

---

**Algorithm 4**

---

- 1: **Parameters:** Stepsizes  $\eta > 0$ , momentums  $\alpha, \gamma$ , batchsize  $b \in \{1, \dots, n\}$ , probability  $p \in (0, 1)$
  - 2: **Initialization:** Choose  $z^0 = w^0 \in \text{dom } g$ . Put  $z^{-1} = z^0, w^{-1} = w^0$
  - 3: **for**  $k = 0, 1, 2, \dots$  **do**
  - 4:   Sample  $j_1^k, \dots, j_b^k$  independently from  $\{1, \dots, m\}$  uniformly at random
  - 5:    $S^k = \{j_1^k, \dots, j_b^k\}$
  - 6:    $\Delta^k = \frac{1}{b} \sum_{j \in S^k} (F_j(x^k) - F_j(w^{k-1}) + \alpha(F_j(x^k) - F_j(x^{k-1}))) + F(w^{k-1})$
  - 7:    $x^{k+1} = \text{prox}_{\eta g}(x^k + \gamma(w^k - x^k) - \eta \Delta^k)$
  - 8:    $w^{k+1} = \begin{cases} x^{k+1}, & \text{with probability } p \\ w^k, & \text{with probability } 1 - p \end{cases}$
  - 9: **end for**
- 

---

**Algorithm 5**

---

- Parameters:** Stepsizes  $\eta, \theta > 0$ , momentums  $\alpha, \beta, \gamma$   
**Initialization:** Choose  $\mathbf{z}^0 = \mathbf{w}^0 \in (\text{dom } g)^M, \mathbf{y}^0 \in \mathcal{L}^\perp$ . Put  $\mathbf{z}^{-1} = \mathbf{z}^0, \mathbf{w}^{-1} = \mathbf{w}^0, \mathbf{y}^{-1} = \mathbf{y}^0$
- for**  $k = 0, 1, 2, \dots$  **do**  
   $\Delta^k = \mathbf{F}(x^k) + \alpha(\mathbf{F}(x^k) - \mathbf{F}(x^{k-1})) - (y^k + \alpha(y^k - y^{k-1}))$   
   $x^{k+1} = \text{prox}_{\eta g}(x^k - \eta \Delta^k)$   
   $y^{k+1} = y^k - \theta(\mathbf{W} \otimes \mathbf{I}_d)(x^{k+1} - \beta(\mathbf{F}(x^{k+1}) - y^k))$   
**end for**
- 

---

**Algorithm 6**

---

- 1: **Parameters:** Stepsizes  $\eta_z, \eta_y, \eta_x, \theta > 0$ , momentums  $\alpha, \gamma, \omega, \tau$ , parameters  $\nu, \beta$
  - 2: **Initialization:** Choose  $\mathbf{z}^0 = \mathbf{w}^0 \in (\text{dom } g)^M, \mathbf{y}^0 \in (\mathbb{R}^d)^M, \mathbf{x}^0 \in \mathcal{L}^\perp$ . Put  $\mathbf{z}^{-1} = \mathbf{z}^0, \mathbf{w}^{-1} = \mathbf{w}^0, \mathbf{y}_f = \mathbf{y}^{-1} = \mathbf{y}^0, \mathbf{x}_f = \mathbf{x}^{-1} = \mathbf{x}^0, m_0 = \mathbf{0}^{dM}$
  - 3: **for**  $k = 0, 1, 2, \dots$  **do**
  - 4:    $\Delta_z^k = \mathbf{F}(\mathbf{z}^k) + \alpha(\mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1})) - \nu \mathbf{z}^k - \mathbf{y}^k - \alpha(\mathbf{y}^k - \mathbf{y}^{k-1})$
  - 5:    $\mathbf{z}^{k+1} = \text{prox}_{\eta_z g}(\mathbf{z}^k - \eta_z \Delta_z^k)$
  - 6:    $\mathbf{y}_c^k = \tau \mathbf{y}^k + (1 - \tau) \mathbf{y}_f^k$
  - 7:    $\mathbf{x}_c^k = \tau \mathbf{x}^k + (1 - \tau) \mathbf{x}_f^k$
  - 8:    $\Delta_y^k = \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \mathbf{z}^{k+1} + \gamma(\mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k)$
  - 9:    $\Delta_x^k = \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \beta(\mathbf{x}^k + \mathbf{F}(\mathbf{z}^k))$
  - 10:    $\mathbf{y}^{k+1} = \mathbf{y}^k - \eta_y \Delta_y^k$
  - 11:    $\mathbf{x}^{k+1} = \mathbf{x}^k - (\mathbf{W}_T(Tk) \otimes \mathbf{I}_d)(\eta_x \Delta_x^k + m^k)$
  - 12:    $m^{k+1} = \eta_x \Delta_x^k + m^k - (\mathbf{W}_T(Tk) \otimes \mathbf{I}_d)(\eta_x \Delta_x^k + m^k)$
  - 13:    $\mathbf{y}_f^{k+1} = \mathbf{y}_c^k + \tau(\mathbf{y}^{k+1} - \mathbf{y}^k)$
  - 14:    $\mathbf{x}_f^{k+1} = \mathbf{x}_c^k - \theta(\mathbf{W}_T(Tk) \otimes \mathbf{I}_d)(\mathbf{y}_c^k + \mathbf{x}_c^k)$
  - 15: **end for**
-

## C Proof of Theorems 3.2 and 3.4

The idea of obtaining lower bounds is to find some "bad" example in the class of problems. In particular, we consider saddle point problems (as a special case of VIs) and look for the "bad" function among them. The paper [72] gives such an example for non-distributed deterministic problems. To get a distributed bounds, this problem needs to be divided between computing devices, which are connected in some kind of "bad" computing network [63, 39]. Then on each of the devices the function must be further divided in order to obtain a stochastic sum type problem [30, 29].

As stated above, to obtain lower bounds, we consider a particular case of the variational inequality (1)+(5), the saddle point problem:

$$\min_{x \in \mathbb{R}^d} \max_{y \in \mathbb{R}^d} f(x, y) := \sum_{m=1}^M f_m(x, y) := \sum_{m=1}^M \frac{1}{n} \sum_{i=1}^n f_{m,i}(x, y). \quad (15)$$

In this case  $F(z) = F(x, y) = [\nabla_x f(x, y), -\nabla_y f(x, y)]$  and  $g \equiv 0$ . Next we rewrite Assumptions 2.1 and 2.2 for (15).

**Assumption C.1.** Suppose that

- each  $f_m$  is  $\bar{L}$ -average smooth, on  $\mathbb{R}^d \times \mathbb{R}^d$ , i.e. for all  $x_1, x_2 \in \mathbb{R}^d, y_1, y_2 \in \mathbb{R}^d$  it holds that

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n (\|\nabla_x f_{m,i}(x_1, y_1) - \nabla_x f_{m,i}(x_2, y_2)\|^2 + \|\nabla_y f_{m,i}(x_1, y_1) - \nabla_y f_{m,i}(x_2, y_2)\|^2) \\ \leq \bar{L}^2 (\|x_1 - x_2\|^2 + \|y_1 - y_2\|^2); \end{aligned} \quad (16)$$

- each  $f_m$  is  $L$ -smooth, on  $\mathbb{R}^d \times \mathbb{R}^d$ , i.e. for all  $x_1, x_2 \in \mathbb{R}^d, y_1, y_2 \in \mathbb{R}^d$  it holds that

$$\begin{aligned} \|\nabla_x f_m(x_1, y_1) - \nabla_x f_m(x_2, y_2)\|^2 + \|\nabla_y f_m(x_1, y_1) - \nabla_y f_m(x_2, y_2)\|^2 \\ \leq L^2 (\|x_1 - x_2\|^2 + \|y_1 - y_2\|^2); \end{aligned} \quad (17)$$

- $f$  is  $\mu$ -strongly-convex-strongly-concave on  $\mathbb{R}^d \times \mathbb{R}^d$ , i.e. for all  $x_1, x_2 \in \mathbb{R}^d, y_1, y_2 \in \mathbb{R}^d$  it holds that

$$\begin{aligned} \langle \nabla_x f(x_1, y_1) - \nabla_x f(x_2, y_2); x_1 - x_2 \rangle - \langle \nabla_y f(x_1, y_1) - \nabla_y f(x_2, y_2); y_1 - y_2 \rangle \\ \geq \mu (\|x_1 - x_2\|^2 + \|y_1 - y_2\|^2). \end{aligned} \quad (18)$$

Next we rewrite Definition 3.1 for (15). Since  $g \equiv 0$ , then  $\text{prox}_{\rho g}(z) = z$ .

**Definition C.2** (Oracle). Each agent  $m$  has its own local memories  $\mathcal{M}_m^x$  and  $\mathcal{M}_m^y$  for the  $x$ - and  $y$ -variables, respectively—with initialization  $\mathcal{M}_m^x = \mathcal{M}_m^y = \{0\}$ .  $\mathcal{M}_m^x$  and  $\mathcal{M}_m^y$  are updated as follows.

- **Local computation:** At each local iteration device  $m$  can sample uniformly and independently batch  $S_m$  of any size  $b$  from  $\{f_{m,i}\}$  and adds to its  $\mathcal{M}_m^x$  and  $\mathcal{M}_m^y$  a finite number of points  $x, y$ , satisfying

$$\begin{aligned} x \in \text{span}\left\{x', \sum_{i_m \in S_m} \nabla_x f_{m,i_m}(x'', y'')\right\}, \\ y \in \text{span}\left\{y', \sum_{i_m \in S_m} \nabla_y f_{m,i_m}(x'', y'')\right\}, \end{aligned} \quad (19)$$

for given  $x', x'' \in \mathcal{M}_m^x$  and  $y', y'' \in \mathcal{M}_m^y$ . Such call needs  $b$  local computations to collect the batch. Batch of the size  $n$  is equal to the full  $f_m$ ;

- **Communication:** Based upon communication rounds among neighbouring nodes, at the communication with the number  $t$ ,  $\mathcal{M}_m^x$  and  $\mathcal{M}_m^y$  are updated according to

$$\mathcal{M}_m^x := \text{span}\left\{\bigcup_{(i,m) \in \mathcal{E}(t)} \mathcal{M}_i^x\right\}, \quad \mathcal{M}_m^y := \text{span}\left\{\bigcup_{(i,m) \in \mathcal{E}(t)} \mathcal{M}_i^y\right\}. \quad (20)$$

- **Output:** The final global output is calculated as:

$$\hat{x} \in \text{span}\left\{\bigcup_{m=1}^M \mathcal{M}_m^x\right\}, \quad \hat{y} \in \text{span}\left\{\bigcup_{m=1}^M \mathcal{M}_m^y\right\}.$$

We construct the following bilinearly functions with  $\bar{L}, \mu$ . Let us consider a communication graph  $G$  with vertexes  $\{1, \dots, M\}$ . Define  $B = \{1\}$  and  $\bar{B} = \{M\}$ , with  $|B| = |\bar{B}| = 1$ . We then construct the following



$$\begin{aligned}
&= \frac{\bar{L}^2}{8} \sum_{i=1}^n (y_1 - y_2)^T \left[ \sum_{j \equiv (i-1) \pmod n} \left( a_{1,2j+1} e_{2j+1}^T + a_{1,2j+2} e_{2j+2}^T \right) \right] \\
&\quad \left[ \sum_{j \equiv (i-1) \pmod n} \left( e_{2j+1} a_{1,2j+1}^T + e_{2j+2} a_{1,2j+2}^T \right) \right] (y_1 - y_2) \\
&+ \frac{\bar{L}^2}{8} \sum_{i=1}^n (x_1 - x_2)^T \left[ \sum_{j \equiv (i-1) \pmod n} \left( e_{2j+1} a_{1,2j+1}^T + e_{2j+2} a_{1,2j+2}^T \right) \right] \\
&\quad \left[ \sum_{j \equiv (i-1) \pmod n} \left( a_{1,2j+1} e_{2j+1}^T + a_{1,2j+2} e_{2j+2}^T \right) \right] (x_1 - x_2) \\
&+ \frac{2\mu^2}{9} \|x_1 - x_2\|^2 + \frac{2\mu^2}{9} \|y_1 - y_2\|^2 \\
&= \frac{\bar{L}^2}{8} \sum_{i=1}^n (y_1 - y_2)^T \left[ \sum_{j \equiv (i-1) \pmod n} \left( a_{1,2j+1} a_{1,2j+1}^T + a_{1,2j+2} a_{1,2j+2}^T \right) \right] (y_1 - y_2) \\
&+ \frac{\bar{L}^2}{8} \sum_{i=1}^n (x_1 - x_2)^T \left[ \sum_{j \equiv (i-1) \pmod n} \left( e_{2j+1} e_{2j+1}^T + 5e_{2j+2} e_{2j+2}^T \right) \right] (x_1 - x_2) \\
&+ \frac{2\mu^2}{9} \|x_1 - x_2\|^2 + \frac{2\mu^2}{9} \|y_1 - y_2\|^2 \\
&\leq \frac{\bar{L}^2}{8} \sum_{j=1}^d (y_1 - y_2)^T \left( a_{1,j} a_{1,j}^T \right) (y_1 - y_2) + \frac{5\bar{L}^2}{8} \sum_{j=1}^d (x_1 - x_2)^T \left( e_j e_j^T \right) (x_1 - x_2) \\
&\quad + \frac{2\mu^2}{9} \|x_1 - x_2\|^2 + \frac{2\mu^2}{9} \|y_1 - y_2\|^2 \\
&\leq \frac{\bar{L}^2}{8} \sum_{j=1}^d (y_1 - y_2)^T \left( a_{1,j} a_{1,j}^T \right) (y_1 - y_2) + \frac{5\bar{L}^2}{8} \|x_1 - x_2\|^2 \\
&\quad + \frac{2\mu^2}{9} \|x_1 - x_2\|^2 + \frac{2\mu^2}{9} \|y_1 - y_2\|^2 \\
&\leq \frac{3\bar{L}^2}{4} \|y_1 - y_2\|^2 + \frac{5\bar{L}^2}{8} \|x_1 - x_2\|^2 + \frac{2\mu^2}{9} \|x_1 - x_2\|^2 + \frac{2\mu^2}{9} \|y_1 - y_2\|^2.
\end{aligned}$$

The last inequality follows from  $\lambda_{\max} \left( \sum_{j=1}^d (a_{1,j} a_{1,j}^T) \right) \leq 6$ . Finally, with  $\mu \leq \bar{L}$  we get

$$\begin{aligned}
&\frac{1}{n} \sum_{i=1}^n \left[ \|\nabla_x f_{1,i}(x_1, y_1) - \nabla_x f_{1,i}(x_2, y_2)\|^2 + \|\nabla_y f_{1,i}(x_1, y_1) - \nabla_y f_{1,i}(x_2, y_2)\|^2 \right] \\
&\leq \bar{L}^2 (\|x_1 - x_2\|^2 + \|y_1 - y_2\|^2).
\end{aligned}$$

□

The next two lemmas give an idea of how quickly we approximate the solution of (23) depending on the number of communications and local iterations. For simplicity, we divide a situation in two parts: one part is devoted to communications (taking into account the fact that we are not limited in the number of local iterations), the second - on the contrary (we concentrate on local computations and assume that communications cost nothing).

**Lemma C.4.** *Let Problem (21) be solved by any method that satisfies Definition C.2. Then after  $K$  communication rounds, only the first  $\lfloor \frac{K}{\bar{L}} \rfloor$  coordinates of the global output can be non-zero while the rest of the  $d - \lfloor \frac{K}{\bar{L}} \rfloor$  coordinates are strictly equal to zero. Here  $l$  is "distance" between  $B$  and  $\bar{B}$  (how quickly can we transfer information from  $B$  to  $\bar{B}$ ).*

*Proof.* We begin introducing some notation for our proof. Let

$$E_0 := \{0\}, \quad E_K := \text{span}\{e_1, \dots, e_K\}.$$

Note that, the initialization gives  $\mathcal{M}_n^x = E_0, \mathcal{M}_m^y = E_0$ .

Suppose that, for some  $m$ ,  $\mathcal{M}_m^x = E_K$  and  $\mathcal{M}_m^y = E_K$ , at some given time. Let us analyze how  $\mathcal{M}_m^x, \mathcal{M}_m^y$  can change by performing only local computations.

Firstly, we consider the case when  $K$  odd. After one local update, we have the following:

- For machines  $m$  which own  $f_1$ , it holds

$$\begin{aligned} x &\in \text{span}\{e_1, x', A_1 y'\} = E_K, \\ y &\in \text{span}\{e_1, y', A_1^T x'\} = E_K, \end{aligned} \tag{24}$$

for given  $x' \in \mathcal{M}_m^x$  and  $y' \in \mathcal{M}_m^y$ . In details, each local iteration uses matrices  $A_{1,i}$  (in the stochastic) or  $A_1$  (in the deterministic). But here we talk only about communications and do not pay attention to the number of local iterations. Therefore, without loss of generality, we can immediately assume that all local calculations change our output according to (24). Since  $A_1$  has a block diagonal structure, after local computations, we have  $\mathcal{M}_m^x = E_K$  and  $\mathcal{M}_m^y = E_K$ . The situation does not change, no matter how many local computations one does.

- For machines  $m$  which own  $f_2$ , it holds

$$\begin{aligned} x &\in \text{span}\{x', A_2 y'\} = E_{K+1}, \\ y &\in \text{span}\{y', A_2^T x'\} = E_{K+1}, \end{aligned}$$

for given  $x' \in \mathcal{M}_m^x$  and  $y' \in \mathcal{M}_m^y$ . It means that, after local computations, one has  $\mathcal{M}_m^x = E_{K+1}$  and  $\mathcal{M}_m^y = E_{K+1}$ . Therefore, machines with function  $f_2$  can progress by one new non-zero coordinate.

This means that we constantly have to transfer progress from the group of machines with  $f_1$  to the group of machines with  $f_2$  and back. Initially, all devices have zero coordinates. Further, machines with  $f_1$  can receive the first nonzero coordinate (but only the first, the second is not), and the rest of the devices are left with all zeros. Next, we pass the first non-zero coordinate to machines with  $f_2$ . To do this,  $l$  communication rounds are needed. By doing so, they can make the second coordinate non-zero, and then transfer this progress to the machines with  $f_1$ . Then the process continues in the same way. This completes the proof.  $\square$

In the next lemma, we will give an understanding of how local progress towards a solution occurs. For this, we will assume that communications cost nothing.

**Lemma C.5.** *Let Problem (21) be solved by any method that satisfies Definition C.2. Then after  $N$  local calls (for each node), in expectation only the first  $\lfloor \frac{N}{n} \rfloor$  coordinates of the global output can be non-zero while the rest of the  $d - \lfloor \frac{N}{n} \rfloor$  coordinates are strictly equal to zero.*

*Proof.* As is clear from the previous lemma, communications make sense if an update ( $E_K \rightarrow E_{K+1}$ ) is reached on one of the nodes. Depending on  $K$ , this happens on the nodes with  $f_1$  or  $f_2$  (but not simultaneously). The question is how many local calls should be made to get this update. One can understand it by looking at the structure of matrices  $A_{1,i}$  and  $A_{2,i}$  from (22). Only one of  $n$  matrices is suitable for us. For example, in case of  $K = 2k$ , we need  $f_{1,j}$  with  $j \equiv k \pmod n$ .

Suppose that  $s_1, s_2, \dots$  times call stochastic oracle with batchsize  $1, 2, \dots$ . Then  $\sum_{j=1}^n j s_j + N$ . Due to the fact that the choice of batches is random and uniform, the random variable responsible for the total number of updates during the operation of the algorithm has the sum of binomial distribution with pairs of parameters  $(s_j; \frac{j}{n})$ . It remains only to take the mathematical expectation and the lemma is proved.  $\square$

The next lemma is devoted to provide an approximate solution of problem (23), and shows that this approximation is close to a real solution. The proof of the lemma follows closely that of Lemma 3.3 from [72], and is reported for the sake of completeness.

**Lemma C.6** (Lemma 3.3 from [72]). *Let  $\alpha = \frac{2n\mu^2}{L^2}$  and  $q = \frac{1}{2}(2 + \alpha - \sqrt{\alpha^2 + 4\alpha}) \in (0; 1)$ —the smallest root of  $q^2 - (2 + \alpha)q + 1 = 0$ ; and let define*

$$\bar{y}_i^* = \frac{q^i}{1 - q}, \quad i \in [d].$$

*The following bound holds when  $\bar{y}^* := [y_1^*, \dots, y_d^*]^\top$  is used to approximate the solution  $y^*$ :*

$$\|\bar{y}^* - y^*\| \leq \frac{q^{d+1}}{\alpha(1 - q)}.$$

*Proof.* Let us write the dual function for (23):

$$h(y) = -\frac{1}{2}y^T \left( \frac{L^2}{2n\mu} A^T A + \mu I \right) y + \frac{L^2}{2n\mu} e_1^T y,$$



$\frac{1}{2}(2 + \alpha - \sqrt{\alpha^2 + 4\alpha}) \in (0; 1)$ . Then, any output  $\hat{x}, \hat{y}$  produced by any method satisfying Definition C.2 after  $K$  communications and  $N$  local calls, is such that

$$\mathbb{E} [\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2] \geq \left( q^{\frac{2K}{l}} + q^{\frac{2N}{n}} \right) \frac{\|y_0 - y^*\|^2}{16}.$$

*Proof.* Let us assume that in output we have  $k$  non-zero coordinates. By definition of  $\bar{y}^*$ , with  $q < 1$  and  $k \leq \frac{d}{2}$ , we have

$$\begin{aligned} \|\hat{y} - \bar{y}^*\|^2 &\geq \sqrt{\sum_{j=k+1}^d (\bar{y}_j^*)^2} = \frac{q^k}{1-q} \sqrt{q^2 + q^4 + \dots + q^{2(d-k)}} \\ &\geq \frac{q^k}{\sqrt{2}(1-q)} \sqrt{q^2 + q^4 + \dots + q^{2d}} = \frac{q^k}{\sqrt{2}} \|\bar{y}^*\|^2 = \frac{q^k}{\sqrt{2}} \|y_0 - \bar{y}^*\|^2. \end{aligned}$$

Using Lemma C.6 for  $d \geq 2 \log_q \left( \frac{\alpha}{4\sqrt{2}} \right)$  we can guarantee that  $\bar{y}^* \approx y^*$  (for more detailed proof see [72]) and

$$\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2 \geq \|\hat{y} - y^*\|^2 \geq \frac{q^{2k}}{16} \|y_0 - y^*\|^2.$$

It remains only to note that  $k$  depends on the number of nonzero coordinates from communications  $k_c$  and local computations  $k_l$ . For this we use Lemmas C.4 and C.5:

$$\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2 \geq \frac{q^{2k}}{16} \|y_0 - y^*\|^2 \geq \frac{q^{2 \min(k_c, k_l)}}{16} \|y_0 - y^*\|^2 \geq \frac{q^{k_c + k_l}}{16} \|y_0 - y^*\|^2.$$

By Lemma C.4 we have  $k_c \leq \lfloor \frac{K}{l} \rfloor$ , where  $l$  is "distance" between  $B$  and  $\bar{B}$ . By Lemma C.5 we get that  $k_l$  has binomial distribution with parameters  $N$  and  $\frac{1}{n}$ .

$$\begin{aligned} \mathbb{E} [\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2] &\geq \frac{q^{2 \lfloor \frac{K}{l} \rfloor}}{32} \|y_0 - y^*\|^2 + \mathbb{E} \left[ \frac{q^{2k_l}}{32} \|y_0 - y^*\|^2 \right] \\ &\geq \frac{q^{\frac{2K}{l}}}{32} \|y_0 - y^*\|^2 + \frac{\mathbb{E} [q^{2k_l}]}{32} \|y_0 - y^*\|^2 \\ &\geq \left( q^{\frac{2K}{l}} + q^{2\mathbb{E}[k_l]} \right) \cdot \frac{1}{32} \|y_0 - y^*\|^2 \\ &\geq \left( q^{\frac{2K}{l}} + q^{\frac{2N}{n}} \right) \cdot \frac{1}{32} \|y_0 - y^*\|^2. \end{aligned}$$

Here we use Jensen's inequality. □

It remains to get an estimate on  $l$  ("distance" between  $B$  and  $\bar{B}$ ). For the fixed network it is real distance, for time-varying – rate how fast information transmits in the network from  $B$  to  $\bar{B}$ .

## C.1 Fixed network

**Theorem C.8** (Theorem 3.2). *Let  $\bar{L} > \mu > 0$ ,  $n \in \mathbb{N}$  (with  $\bar{L}/\mu \geq \sqrt{n}$ ),  $\chi \geq 1$  and  $K, N \in \mathbb{N}$ . There exists a distributed saddle-point problem over fixed network (Assumption 2.3). For which the following statements are true:*

- a gossip matrix  $\mathbf{W}$  has  $\chi(\mathbf{W}) = \chi$ ,
- $f = \sum_{m=1}^M \frac{1}{n} \sum_{i=1}^n f_{m,i} : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is  $\mu$ -strongly-convex-strongly-concave,
- $f_m$  are  $\bar{L}$ -average smooth and  $L = \frac{\bar{L}}{\sqrt{n}}$ -smooth,
- size  $d \geq \max \left\{ 2 \log_q \left( \frac{\alpha}{4\sqrt{2}} \right), 2K, 2N \right\}$ , where  $\alpha = \frac{2n\mu^2}{L^2}$  and  $q = \frac{1}{2} (2 + \alpha - \sqrt{\alpha^2 + 4\alpha}) \in (0; 1)$ ,
- the solution of the problem is non-zero:  $x^* \neq 0, y^* \neq 0$ .

Then for any output  $\hat{z}$  of any procedure (Definition C.2) with  $K$  communication rounds and  $N$  local computations, one can obtain the following estimate:

$$\|\hat{z} - z^*\|^2 = \Omega \left( \exp \left( -\frac{80}{1 + \sqrt{\frac{2L^2}{\mu^2} + 1}} \cdot \frac{K}{\sqrt{\chi}} \right) \|y_0 - y^*\|^2 \right);$$



$$\|\hat{z} - z^*\|^2 = \Omega \left( \exp \left( -\frac{16}{n + \sqrt{\frac{2n\bar{L}^2}{\mu^2} + n^2}} \cdot N \right) \|y_0 - y^*\|^2 \right).$$

*Proof.* Applying Lemma C.7, we have

$$\left(\frac{1}{q}\right)^{\frac{2K}{l}} \geq \frac{\|y_0 - y^*\|^2}{32\mathbb{E}[\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]} \quad \text{and} \quad \left(\frac{1}{q}\right)^{\frac{2N}{n}} \geq \frac{\|y_0 - y^*\|^2}{32\mathbb{E}[\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]}.$$

Taking the logarithm on both sides, we get

$$\frac{2K}{l} \geq \ln \left( \frac{\|y_0 - y^*\|^2}{32\mathbb{E}[\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]} \right) \frac{1}{\ln(q^{-1})}.$$

Next, we work with

$$\begin{aligned} \frac{1}{\ln(q^{-1})} &= \frac{1}{\ln(1 + (1-q)/q)} \geq \frac{q}{1-q} = \frac{1 + \frac{n\mu^2}{\bar{L}^2} - \sqrt{\frac{2n\mu^2}{\bar{L}^2} + \left(\frac{n\mu^2}{\bar{L}^2}\right)^2}}{\sqrt{\frac{2n\mu^2}{\bar{L}^2} + \left(\frac{n\mu^2}{\bar{L}^2}\right)^2} - \frac{n\mu^2}{\bar{L}^2}} \\ &\geq \frac{\sqrt{\frac{2n\mu^2}{\bar{L}^2} + \left(\frac{n\mu^2}{\bar{L}^2}\right)^2} + \frac{n\mu^2}{\bar{L}^2}}{\frac{8n\mu^2}{\bar{L}^2}} = \frac{1}{8} \left( 1 + \sqrt{\frac{2\bar{L}^2}{n\mu^2} + 1} \right). \end{aligned}$$

One can then obtain

$$\frac{2K}{l} \geq \ln \left( \frac{\|y_0 - y^*\|^2}{32\mathbb{E}[\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]} \right) \cdot \frac{1}{8} \left( 1 + \sqrt{\frac{2\bar{L}^2}{n\mu^2} + 1} \right),$$

and

$$\exp \left( \frac{1}{1 + \sqrt{\frac{2\bar{L}^2}{n\mu^2} + 1}} \frac{16K}{l} \right) \geq \frac{\|y_0 - y^*\|^2}{32\mathbb{E}[\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]}.$$

The next proof follow similar steps as in the proof of Theorem 2 from [63]. Let  $\gamma_M = \frac{1 - \cos \frac{\pi}{M}}{1 + \cos \frac{\pi}{M}}$  be a decreasing sequence of positive numbers. Since  $\gamma_2 = 1$  and  $\lim_m \gamma_M = 0$ , there exists  $M \geq 2$  such that  $\gamma_M \geq \chi^{-1} > \gamma_{M+1}$ .

• If  $M \geq 3$ , let us consider linear graph of size  $M$  with vertexes  $v_1, \dots, v_M$ , and weighted with  $w_{1,2} = 1 - a$  and  $w_{i,i+1} = 1$  for  $i \geq 2$ . Then we applied Lemmas 1 and 3 and get:

$$\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2 \geq q \frac{2K}{l} \frac{\|y_0 - y^*\|^2}{32}.$$

If  $\mathbf{W}_a$  is the normalized Laplacian of the weighted graph  $\mathcal{G}$ , one can note that with  $a = 0$ ,  $\chi^{-1}(\mathbf{W}_a) = \gamma_M$ , with  $a = 1 - \chi^{-1}(\mathbf{W}_a) = 0$ . Hence, there exists  $a \in (0; 1]$  such that  $\chi^{-1}(\mathbf{W}_a) = \chi^{-1}$ . Then  $\chi^{-1} \geq \gamma_{M+1} \geq \frac{2}{(M+1)^2}$ , and  $M \geq \sqrt{2\chi} - 1 \geq \frac{\sqrt{\chi}}{4}$ . Finally,  $l = M - 1 \geq \frac{15M}{16} - 1 \geq \frac{15}{16} (\sqrt{2\chi} - 1) - 1 \geq \frac{\sqrt{\chi}}{5}$  since  $\chi^{-1} \leq \gamma_3 = \frac{1}{3}$ . Hence,

$$\exp \left( 80 \frac{K}{\sqrt{\chi}} \cdot \frac{1}{1 + \sqrt{\frac{2\bar{L}^2}{n\mu^2} + 1}} \right) \geq \frac{\|y_0 - y^*\|^2}{32(\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2)}. \quad (25)$$

• If  $M = 2$ , we construct a totally connected network with 3 nodes with weight  $w_{1,3} = a \in [0; 1]$ . Let  $\mathbf{W}_a$  is the normalized Laplacian. If  $a = 0$ , then the network is a linear graph and  $\chi^{-1}(\mathbf{W}_a) = \gamma_3 = \frac{1}{3}$ . Hence, there exists  $a \in [0; 1]$  such that  $\chi^{-1}(\mathbf{W}_a) = \chi^{-1}$ . Finally,  $B = \{v_1\}$ ,  $\bar{B} = \{v_3\}$  and  $l \geq 1 \geq \frac{1}{2\sqrt{\chi^{-1}}}$ . Whence it follows that in this case (25) is also valid.

The same way we can work with (but without considering graph):

$$\left(\frac{1}{q}\right)^{\frac{2N}{n}} \geq \frac{\|y_0 - y^*\|^2}{32\mathbb{E}[\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]}.$$

□

## C.2 Time-varying network

**Theorem C.9** (Theorem 3.4). *Let  $\bar{L} > \mu > 0$ ,  $n \in \mathbb{N}$  (with  $\bar{L}/\mu \geq \sqrt{n}$ ),  $\hat{\chi} \geq 3$  and  $K, N \in \mathbb{N}$ . There exists a distributed saddle-point problem over time-varying network (Assumption 2.4). For which the following statements are true:*

- Assumption 2.4 holds with  $\chi = \hat{\chi}$ ,
- $f = \sum_{m=1}^M \frac{1}{n} \sum_{i=1}^n f_{m,i} : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$  is  $\mu$ -strongly-convex-strongly-concave,
- $f_m$  are  $\bar{L}$ -average smooth and  $L = \frac{\bar{L}}{\sqrt{n}}$ -smooth,
- size  $d \geq \max \left\{ 2 \log_q \left( \frac{\alpha}{4\sqrt{2}} \right), 2K, 2N \right\}$ , where  $\alpha = \frac{2n\mu^2}{L^2}$  and  $q = \frac{1}{2} (2 + \alpha - \sqrt{\alpha^2 + 4\alpha}) \in (0; 1)$ ,
- the solution of the problem is non-zero:  $x^* \neq 0, y^* \neq 0$ .

Then for any output  $\hat{z}$  of any procedure (Definition C.2) with  $K$  communication rounds and  $N$  local computations, one can obtain the following estimate:

$$\|\hat{z} - z^*\|^2 = \Omega \left( \exp \left( -\frac{64}{1 + \sqrt{\frac{2\bar{L}^2}{\mu^2} + 1}} \cdot \frac{K}{B\hat{\chi}} \right) \|y_0 - y^*\|^2 \right);$$

$$\|\hat{z} - z^*\|^2 = \Omega \left( \exp \left( -\frac{16}{n + \sqrt{\frac{2n\bar{L}^2}{\mu^2} + n^2}} \cdot N \right) \|y_0 - y^*\|^2 \right).$$

*Proof.* The same way as in the previous Theorem we can obtain

$$\exp \left( \frac{1}{1 + \sqrt{\frac{2\bar{L}^2}{n\mu^2} + 1}} \frac{16K}{l} \right) \geq \frac{\|y_0 - y^*\|^2}{32\mathbb{E} [\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]}.$$

Following [39], we can consider the next sequences of graphs. Let us choose  $M = \lfloor \hat{\chi} \rfloor$ . Each communication  $t$  such that  $t \neq 8\tau$  (for any  $\tau \in \mathbb{N}$ ) we consider empty network. Each communication  $t$  such that  $t = 8\tau$  (for some  $\tau \in \mathbb{N}$ ) we construct star graph with vertex  $[(\tau - 1) \bmod (M - 2)] + 2$  in the center of this star. It means that in the 8th communication vertex 2 is in the center; in the 16th communication vertex 3 is in the center etc. One can note that only vertexes with  $f_3$  are in the center and they change sequentially. As matrices  $\mathbf{W}(t)$  we consider the normalized Laplacians. Then it holds  $\chi = M = \lfloor \hat{\chi} \rfloor$ . It holds that Assumption 2.4 is valid with  $\hat{\chi}$ .

It is left to estimate  $l$ . Suppose we need to transfer information from  $B$  to  $\bar{B}$ . At best, the following will happen:

- vertex  $j$  in the center, it means that we can transfer information to  $j$ ;
- $B - 1$  empty graphs - "empty" communications;
- vertex  $j + 1$  in the center, it means that we can transfer information to  $j + 1$ ;
- ...
- vertex  $j - 1$  in the center, it means that we can transfer information to  $j - 1$  (still there is no information in  $\bar{B}$ );
- $B - 1$  empty graphs - "empty" communications;
- vertex  $j$  in the center, now we can transfer from  $j$  to  $\bar{B}$ .

In this (the best variant) we spend  $M - 1 + (B - 1)(M - 2)$  communication rounds. It means that  $l \geq M - 1 + (B - 1)(M - 2) \geq B(M - 2) = B(\lfloor \hat{\chi} \rfloor - 2) \geq \frac{B\hat{\chi}}{4}$  (for  $\hat{\chi} \geq 3$ ). Then we get

$$\exp \left( \frac{1}{1 + \sqrt{\frac{2\bar{L}^2}{n\mu^2} + 1}} \frac{64K}{B\hat{\chi}} \right) \geq \frac{\|y_0 - y^*\|^2}{32\mathbb{E} [\|\hat{x} - x^*\|^2 + \|\hat{y} - y^*\|^2]}.$$

□

## D Proof of Theorem 4.1

We start the proof from the following lemma on  $\delta^k$  and  $\Delta^{k+1/2}$  from Algorithm 1.

**Lemma D.1.** *The following inequality holds:*

$$\mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] \leq \frac{2\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 + \alpha^2 \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right]. \quad (26)$$

$$\mathbb{E}_k \left[ \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \right] \leq \frac{2\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + 2L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right]. \quad (27)$$

where  $\mathbb{E}_k [\delta^k]$  is equal to

$$\mathbb{E}_k [\delta^k] = F(\mathbf{z}^k) + \alpha(F(\mathbf{z}^k) - F(\mathbf{z}^{k-1})). \quad (28)$$

*Proof.* Due to the fact that the batch  $S^k$  is generated uniformly and independently for all workers, we can make sure that (28) is correct. Then using definition of  $\Delta^k$  from Algorithm 1, we get

$$\begin{aligned} \mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] &\leq 2\mathbb{E}_k \left[ \left\| \frac{1}{b} \sum_{j \in S^k} (\mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1})) - (\mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{w}^{k-1})) \right\|^2 \right] \\ &\quad + 2\mathbb{E}_k \left[ \left\| \frac{\alpha}{b} \sum_{j \in S^k} (\mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1})) - (\mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1})) \right\|^2 \right] \end{aligned}$$

By randomness and independence of indexes in  $S^k$ , we obtain

$$\begin{aligned} \mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] &= \frac{2}{b^2} \mathbb{E}_k \left[ \sum_{j \in S^k} \left\| (\mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1})) - (\mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{w}^{k-1})) \right\|^2 \right] \\ &\quad + \frac{2\alpha^2}{b^2} \mathbb{E}_k \left[ \sum_{j \in S^k} \left\| (\mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1})) - (\mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1})) \right\|^2 \right] \end{aligned}$$

The property of the second moment  $\mathbb{E} \|\xi - \mathbb{E}\xi\|^2 = \mathbb{E}\|\xi\|^2 - \|\mathbb{E}\xi\|^2$  gives

$$\begin{aligned} \mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] &\leq \frac{2}{b^2} \mathbb{E}_k \left[ \sum_{j \in S^k} \left\| \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1}) \right\|^2 \right] \\ &\quad + \frac{2\alpha^2}{b^2} \mathbb{E}_k \left[ \sum_{j \in S^k} \left\| \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1}) \right\|^2 \right] \end{aligned}$$

Again from the fact that  $S^k$  is generated uniformly and independently for all samples and workers, we can obtain that for each worker  $m$  indexes  $j_{m,1}^k, \dots, j_{m,b}^k$  have the same uniform distribution, that means

$$\begin{aligned} \mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] &\leq \frac{2}{b^2} \mathbb{E}_k \left[ \sum_{j \in S^k} \left\| \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1}) \right\|^2 \right] \\ &\quad + \frac{2\alpha^2}{b^2} \mathbb{E}_k \left[ \sum_{j \in S^k} \left\| \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1}) \right\|^2 \right] \\ &= \frac{2}{mb} \sum_{j=1}^m \left( \left\| \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{w}^{k-1}) \right\|^2 + \alpha^2 \left\| \mathbf{F}_j(\mathbf{z}^k) - \mathbf{F}_j(\mathbf{z}^{k-1}) \right\|^2 \right). \end{aligned}$$

Using Assumption 2.1, we get

$$\mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] \leq \frac{2\bar{L}^2}{b} \left( \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 + \alpha^2 \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right).$$

This concludes the proof of (26).

The proof chain for (27) is very similar.

$$\mathbb{E} \left[ \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \right] \leq 2\mathbb{E} \left[ \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^{k+1}) \right\|^2 \right] + 2\mathbb{E} \left[ \left\| \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^*) \right\|^2 \right]$$

$$\begin{aligned}
&= \mathbb{E} \left[ \frac{2}{b^2} \mathbb{E}_{k+1/2} \sum_{j \in S^{k+1/2}} \left\| \left( \mathbf{F}_j(\mathbf{z}^{k+1}) - \mathbf{F}_j(\mathbf{w}^k) \right) - \left( \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{w}^k) \right) \right\|^2 \right] \\
&\quad + 2L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\
&\leq \mathbb{E} \left[ \frac{2}{b^2} \mathbb{E}_{k+1/2} \sum_{j \in S^{k+1/2}} \left\| \mathbf{F}_j(\mathbf{z}^{k+1}) - \mathbf{F}_j(\mathbf{w}^k) \right\|^2 \right] + 2L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\
&\leq \frac{2\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + 2L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right].
\end{aligned}$$

Here, we used Assumption 2.1.  $\square$

Before proving the main lemma of this section, let us introduce an auxiliary notation. Throughout the proof, we denote  $\mathbf{W}^\dagger : \text{range} \mathbf{W} \rightarrow \text{range} \mathbf{W}$  the inverse of the map  $\mathbf{W} : \text{range} \mathbf{W} \rightarrow \text{range} \mathbf{W}$ . And we denote  $\|\mathbf{y}\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 = \langle (\mathbf{W}^\dagger \otimes \mathbf{I}_d) \mathbf{y}; \mathbf{y} \rangle$ .

We define the following Lyapunov function:

$$\begin{aligned}
\Psi^k &= \left( \frac{1}{\eta} + \frac{3\mu}{2} \right) \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad + 2 \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle + \frac{1}{2\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
&\quad + \frac{1}{8\eta} \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 + \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 + \frac{\gamma}{2\eta} \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2.
\end{aligned} \tag{29}$$

Here we also use

$$\mathbf{y}^* = \mathbf{P}\mathbf{F}(\mathbf{z}^*) \tag{30}$$

with  $\mathbf{P} \in \mathbb{R}^{nd \times nd}$ , an orthogonal projection matrix onto the subspace  $\mathcal{L}^\perp$ , given as

$$\mathbf{P} = (\mathbf{I}_M - \frac{1}{M} \mathbf{1}_M \mathbf{1}_M^\top) \otimes \mathbf{I}_d,$$

where  $\mathbf{1}_M = (1, \dots, 1)^\top \in \mathbb{R}^M$ .

**Lemma D.2.** Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a fixed connected graph  $\mathcal{G}$  with a gossip matrix  $\mathbf{W}$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 1 with parameters

$$\begin{aligned}
\gamma &\leq \frac{1}{8}, \quad \eta \leq \min \left\{ \frac{\sqrt{\alpha\gamma b}}{\sqrt{8 \cdot \bar{L}}}, \frac{1}{16L} \right\}, \quad \beta \leq \min \left\{ \frac{\mu}{4L^2}; \frac{b\gamma}{4\eta L^2} \right\}, \quad \theta \leq \min \left\{ \frac{1}{2\beta}; \frac{1}{16\eta} \right\}, \\
\alpha &= \max \left[ \left( 1 - \frac{\mu\eta}{4} \right); (1 - \beta\theta\chi^{-1}); \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \right].
\end{aligned}$$

Then, after  $k$  iterations we get

$$\mathbb{E} \left[ \frac{1}{2\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \leq \max \left[ \left( 1 - \frac{\mu\eta}{4} \right); (1 - \beta\theta\chi^{-1}); \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \right]^k \cdot \Psi^0.$$

*Proof. Part 1.* We start the proof from considering update of  $\mathbf{y}^{k+1}$  in Algorithm 1.

$$\begin{aligned}
\frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 &= \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad - 2 \langle (\mathbf{W} \otimes \mathbf{I}_d) (\mathbf{z}^{k+1} - \beta(\Delta^{k+1/2} - \mathbf{y}^k)), (\mathbf{W}^\dagger \otimes \mathbf{I}_d) (\mathbf{y}^{k+1} - \mathbf{y}^*) \rangle.
\end{aligned}$$

Next, we use the fact that  $(\mathbf{W} \otimes \mathbf{I}_d)(\mathbf{W}^\dagger \otimes \mathbf{I}_d) = \mathbf{P}$  and obtain

$$\begin{aligned}
\frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 &= \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad - 2 \langle \mathbf{P} (\mathbf{z}^{k+1} - \beta(\Delta^{k+1/2} - \mathbf{y}^k)), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle.
\end{aligned}$$

One can observe, that  $\mathbf{z}^* \in \mathcal{L}$  and then  $\mathbf{P}\mathbf{z}^* = 0$ . Additionally, using update  $\mathbf{y}^{k+1}$ , we can note that  $\mathbf{y}^k \in \mathcal{L}^\perp$  for all  $k = 0, 1, 2, \dots$ . Hence,

$$\frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 = \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2$$

$$\begin{aligned}
& -2\langle \mathbf{P}(\mathbf{z}^{k+1} - \mathbf{z}^*), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + 2\beta\langle \mathbf{P}(\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
& - 2\beta\langle \mathbf{P}\mathbf{y}^k - \mathbf{y}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
= & \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\langle \mathbf{P}(\mathbf{z}^{k+1} - \mathbf{z}^*), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + 2\beta\langle \mathbf{P}(\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
& - 2\beta\langle \mathbf{P}(\mathbf{y}^k - \mathbf{y}^*), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
= & \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + 2\beta\langle \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
& - 2\beta\langle \mathbf{y}^k - \mathbf{y}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle
\end{aligned}$$

By the simple fact  $\|a + b\|^2 = \|a\|^2 + 2\langle a; b \rangle + \|b\|^2$ , we get

$$\begin{aligned}
\frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 & \leq \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + \beta \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 + \beta \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \\
& - \beta \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 - \beta \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \beta \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
= & \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + \beta \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 - \beta \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \\
& + \beta \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2.
\end{aligned}$$

Using the fact, that  $\lambda_{\min}^+(\mathbf{W}) = \chi^{-1}$  and  $\lambda_{\max}(\mathbf{W}) = 1$ , we can note  $-\lambda_{\min}^+(\mathbf{W}^\dagger) \leq -1$  and  $\chi^{-1}\lambda_{\max}(\mathbf{W}^\dagger) \leq 1$  and then get

$$\begin{aligned}
\frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 & \leq \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
& + \beta \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 - \beta\chi^{-1} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 + \beta \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
= & \left( \frac{1}{\theta} - \beta\chi^{-1} \right) \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 - \left( \frac{1}{\theta} - \beta \right) \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
& - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + \beta \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2.
\end{aligned}$$

Then we take a full expectation and use (27):

$$\begin{aligned}
\mathbb{E} \left[ \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \right] & \leq \left( \frac{1}{\theta} - \beta\chi^{-1} \right) \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{W}^\dagger \otimes \mathbf{I}_d)}^2 \right] - \left( \frac{1}{\theta} - \beta \right) \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\
& - 2\mathbb{E} \left[ \langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \right] + 2\beta L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\
& + \frac{2\beta\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right].
\end{aligned} \tag{31}$$

Here we stop and put aside (31). We will not return to it later.

**Part 2.** Now we work with

$$\begin{aligned}
\frac{1}{\eta} \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 & = \frac{1}{\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{2}{\eta} \langle \mathbf{z}^{k+1} - \mathbf{z}^k, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle - \frac{1}{\eta} \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \\
& = \frac{1}{\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{2\gamma}{\eta} \langle \mathbf{w}^k - \mathbf{z}^k, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle - 2\langle \Delta^k - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \\
& - \frac{1}{\eta} \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \\
& - 2\langle \frac{1}{\eta}(\mathbf{z}^k + \gamma(\mathbf{w}^k - \mathbf{z}^k)) - \eta\Delta^k - \mathbf{z}^{k+1}, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle.
\end{aligned}$$

Optimality condition for (1)+(5) it follows, that

$$-F(z^*) \in \partial g(z^*).$$

Let us define  $\Delta^* \in (\mathbb{R}^d)^M$  as

$$\Delta^* = \frac{1}{M}[F(z^*), \dots, F(z^*)]^T.$$

It is to note that  $-\Delta^* \in M\partial g(z^*)$ . On the other hand

$$\Delta^* = (\mathbf{1}_M \mathbf{1}_M^\top \otimes \mathbf{I}_d) \mathbf{F}(z^*) = M(\mathbf{F}(z^*) - \mathbf{P}\mathbf{F}(z^*)) = M(\mathbf{F}(z^*) - \mathbf{y}^*)$$

This means that  $-(\mathbf{F}(z^*) - \mathbf{y}^*) \in \partial g(z^*)$ . From update for  $\mathbf{z}^{k+1}$  of Algorithm 1 it follows, that

$$\mathbf{z}^k + \gamma(\mathbf{w}^k - \mathbf{z}^k) - \eta\Delta^k - \mathbf{z}^{k+1} \in \partial(\eta g)(\mathbf{z}^{k+1}).$$

Hence, from monotonicity of  $\partial g(\cdot)$  we get

$$\begin{aligned} \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 \right] &\leq \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + \frac{2\gamma}{\eta} \mathbb{E} \left[ \langle \mathbf{w}^k - \mathbf{z}^k, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - 2\mathbb{E} \left[ \langle \Delta^k - (\mathbf{F}(z^*) - \mathbf{y}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] - \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] \\ &= \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + \frac{2\gamma}{\eta} \mathbb{E} \left[ \langle \mathbf{w}^k - \mathbf{z}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - \frac{2\gamma}{\eta} \mathbb{E} \left[ \langle \mathbf{z}^k - \mathbf{z}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] - 2\mathbb{E} \left[ \langle \Delta^k - (\mathbf{F}(z^*) - \mathbf{y}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] \\ &= \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^*\|^2 + \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 - \|\mathbf{z}^{k+1} - \mathbf{w}^k\|^2 \right] \\ &\quad - \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 + \|\mathbf{z}^k - \mathbf{z}^*\|^2 - \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] \\ &\quad - 2\mathbb{E} \left[ \langle \Delta^k - (\mathbf{F}(z^*) - \mathbf{y}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] - \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] \\ &= \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^*\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] \\ &\quad - \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^{k+1}\|^2 \right] - 2\mathbb{E} \left[ \langle \Delta^k - (\mathbf{F}(z^*) - \mathbf{y}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - \frac{1-\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right]. \end{aligned}$$

In previous we also use the simple fact  $\|a+b\|^2 = \|a\|^2 + 2\langle a, b \rangle + \|b\|^2$  twice. Small rearrangement gives

$$\begin{aligned} \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 \right] &\leq \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^*\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] \\ &\quad - \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^{k+1}\|^2 \right] - \frac{1-\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] \\ &\quad - 2\mathbb{E} \left[ \langle \mathbb{E}_k [\delta^k] - (\mathbf{y}^k + \alpha(\mathbf{y}^k - \mathbf{y}^{k-1})) - (\mathbf{F}(z^*) - \mathbf{y}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - 2\mathbb{E} \left[ \langle \delta^k - \mathbb{E}_k [\delta^k], \mathbf{z}^{k+1} - \mathbf{z}^k \rangle \right] - 2\mathbb{E} \left[ \langle \delta^k - \mathbb{E}_k [\delta^k], \mathbf{z}^k - \mathbf{z}^* \rangle \right]. \end{aligned}$$

Using the tower property of expectation we can obtain the following:

$$\begin{aligned} \mathbb{E} \left[ \langle \mathbb{E}_k [\delta^k] - \delta^k, \mathbf{z}^k - \mathbf{z}^* \rangle \right] &= \mathbb{E} \left[ \mathbb{E}_k \left[ \langle \mathbb{E}_k [\delta^k] - \delta^k, \mathbf{z}^k - \mathbf{z}^* \rangle \right] \right] \\ &= \mathbb{E} \left[ \langle \mathbb{E}_k [\mathbb{E}_k [\delta^k] - \delta^k], \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ &= \mathbb{E} \left[ \langle \mathbb{E}_k [\delta^k] - \mathbb{E}_k [\delta^k], \mathbf{z}^k - \mathbf{z}^* \rangle \right] = 0. \end{aligned}$$

Hence, with (28)

$$\frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 \right] \leq \frac{1}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right]$$



$$\begin{aligned}
& + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] + \frac{4\alpha^2\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] \\
& - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] + 2\mathbb{E} \left[ \langle \mathbf{y}^{k+1} - \mathbf{y}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
& - 2\alpha\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\
& - 2\alpha\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^{k+1} - \mathbf{z}^k \rangle \right].
\end{aligned}$$

**Part 3.** Now we are ready to consider (31). We sum up previous expression and (31). After small rearrangement, we have

$$\begin{aligned}
& \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
& \leq \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - (2\mu - 2\beta L^2) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\alpha\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
& - \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2 \right] - \frac{1/2 - \gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] \\
& + \frac{4\alpha^2\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - 2\alpha\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}), \mathbf{z}^{k+1} - \mathbf{z}^k \rangle \right] \\
& + 2\alpha\mathbb{E} \left[ \langle \mathbf{y}^k - \mathbf{y}^{k-1}, \mathbf{z}^{k+1} - \mathbf{z}^k \rangle \right] - \left( \frac{1}{\theta} - \beta \right) \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right].
\end{aligned}$$

With  $\theta \leq \frac{1}{2\beta}$  (or  $\beta \leq \frac{1}{2\theta}$ ) and Young's inequality we get

$$\begin{aligned}
& \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
& \leq \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - (2\mu - 2\beta L^2) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\alpha\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
& - \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2 \right] - \frac{1/2 - \gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] \\
& + \frac{4\alpha^2\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] + \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] + 8\eta\alpha^2 \mathbb{E} \left[ \left\| \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) \right\|^2 \right] \\
& + 2\theta\alpha\mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] + \frac{\alpha}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] - \frac{1}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right].
\end{aligned}$$

By choosing  $\theta \leq \frac{1}{16\eta}$  and  $\alpha \leq 1$

$$\begin{aligned}
& \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
& \leq \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - (2\mu - 2\beta L^2) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - 2\alpha\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
& - \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2 \right] - \frac{1/4 - \gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] \\
& + \frac{4\alpha^2\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] + 8\eta\alpha^2 \mathbb{E} \left[ \left\| \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) \right\|^2 \right] + \frac{\alpha}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\
& - \frac{1}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right].
\end{aligned}$$

With  $L$ -Lipshitzness of  $\mathbf{F}$  (Assumption 2.1) and  $\gamma \leq \frac{1}{8}$ , we have

$$\frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right]$$



$$\begin{aligned}
&\leq \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - (2\mu - 2\beta L^2) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad - 2\alpha \mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
&\quad - \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2 \right] - \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] \\
&\quad + \left( \frac{32\alpha^2\bar{L}^2\eta^2}{b} + 64\eta^2\alpha^2L^2 \right) \cdot \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] + \frac{\alpha}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\
&\quad - \frac{1}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right].
\end{aligned}$$

Small rearrangement gives

$$\begin{aligned}
&\frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
&\quad + \frac{1}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] + \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\
&\leq \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - (2\mu - 2\beta L^2) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad - 2\alpha \mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] + \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
&\quad - \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2 \right] + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] \\
&\quad + \left( \frac{32\alpha^2\bar{L}^2\eta^2}{b} + 64\eta^2\alpha^2L^2 \right) \cdot \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] + \frac{\alpha}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right].
\end{aligned}$$

With our choice of  $\eta \leq \min \left\{ \frac{\sqrt{\alpha\gamma b}}{\sqrt{8}\bar{L}}, \frac{1}{16L} \right\}$  and  $\alpha < 1$ , we get

$$\begin{aligned}
&\frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
&\quad + \frac{1}{2\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 + \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\
&\leq \frac{1}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - (2\mu - 2\beta L^2) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] + \alpha \cdot \frac{1}{2\theta} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\
&\quad + \alpha \cdot \frac{1}{8\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] \\
&\quad + \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\gamma}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
&\quad - \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^{k+1} \right\|^2 \right] + \frac{4\bar{L}^2\eta}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right].
\end{aligned}$$

Now, we add  $\frac{\mu}{2} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right]$  to both sides and use update for  $w^{k+1}$  of Algorithm 1

$$\frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \mathbb{E}_{w^{k+1}} \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right] = \frac{\gamma + \frac{1}{2}\eta\mu}{\eta} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{(\gamma + \frac{1}{2}\eta\mu)(1-p)}{\eta p} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right],$$

and get

$$\begin{aligned}
&\left( \frac{1}{\eta} + \frac{\mu}{2} \right) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{\theta} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
&\quad + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] + \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2\theta} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + \frac{1}{8\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] + \left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^{k+1}\|^2 \right] \\
\leq & \left( \frac{1}{\eta} + \frac{\mu}{2} \right) \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] - \left( \frac{3}{2}\mu - 2\beta L^2 \right) \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 \right] \\
& + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \|\mathbf{y}^k - \mathbf{y}^*\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\
& + \alpha \cdot \frac{1}{2\theta} \mathbb{E} \left[ \|\mathbf{y}^k - \mathbf{y}^{k-1}\|^2 \right] + \alpha \cdot \frac{1}{8\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^{k-1}\|^2 \right] \\
& + \left( 1 - p + \frac{p\gamma}{\gamma + \frac{1}{2}\eta\mu} \right) \frac{(\gamma + \frac{1}{2}\eta\mu)}{\eta p} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^*\|^2 \right] + \frac{4\bar{L}^2}{b} \eta \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{w}^{k-1}\|^2 \right].
\end{aligned}$$

Note that  $\eta \leq \min \left\{ \frac{\sqrt{\alpha\gamma b}}{\sqrt{8}\bar{L}}, \frac{1}{16L} \right\}$ ,  $\beta \leq \min \left\{ \frac{\mu}{4L^2}; \frac{b\gamma}{4\eta L^2} \right\}$ , then we get that

$$\left( \frac{\gamma}{\eta} - \frac{2\beta\bar{L}^2}{b} \right) \geq \frac{\gamma}{2\eta}; \quad - \left( \frac{3}{2}\mu - 2\beta L^2 \right) \leq -\mu; \quad \frac{4\bar{L}^2}{b} \eta \leq \alpha \cdot \frac{\gamma}{2\eta}.$$

Hence, it holds

$$\begin{aligned}
& \left( \frac{1}{\eta} + \frac{3\mu}{2} \right) \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 \right] + \frac{1}{\theta} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] + \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \|\mathbf{w}^{k+1} - \mathbf{z}^*\|^2 \right] \\
& + \frac{1}{2\theta} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + \frac{1}{8\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] + \frac{\gamma}{2\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^{k+1}\|^2 \right] \\
\leq & \left( \frac{1}{\eta} + \frac{\mu}{2} \right) \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \|\mathbf{y}^k - \mathbf{y}^*\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\
& + \alpha \cdot \frac{1}{2\theta} \mathbb{E} \left[ \|\mathbf{y}^k - \mathbf{y}^{k-1}\|^2 \right] + \alpha \cdot \frac{1}{8\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^{k-1}\|^2 \right] \\
& + \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \cdot \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^*\|^2 \right] + \alpha \cdot \frac{\gamma}{2\eta} \mathbb{E} \left[ \|\mathbf{x}^k - \mathbf{w}^{k-1}\|^2 \right].
\end{aligned}$$

With our  $\eta$  we have  $\eta\mu \leq 1$  and then

$$\left( \frac{1}{\eta} + \frac{\mu}{2} \right) \leq \left( \frac{1}{\eta} + \frac{3\mu}{2} \right) \left( 1 - \frac{\mu\eta}{4} \right).$$

It gives

$$\begin{aligned}
& \left( \frac{1}{\eta} + \frac{3\mu}{2} \right) \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^*\|^2 \right] + \frac{1}{\theta} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& + 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k+1}) - (\mathbf{y}^k - \mathbf{y}^{k+1}), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] + \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \|\mathbf{w}^{k+1} - \mathbf{z}^*\|^2 \right] \\
& + \frac{1}{2\theta} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + \frac{1}{8\eta} \mathbb{E} \left[ \|\mathbf{z}^{k+1} - \mathbf{z}^k\|^2 \right] + \frac{\gamma}{2\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^{k+1}\|^2 \right] \\
\leq & \left( 1 - \frac{\mu\eta}{4} \right) \cdot \left( \frac{1}{\eta} + \frac{3\mu}{2} \right) \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^*\|^2 \right] + (1 - \beta\theta\chi^{-1}) \cdot \frac{1}{\theta} \|\mathbf{y}^k - \mathbf{y}^*\|_{(\mathbf{w}^\dagger \otimes \mathbf{I}_d)}^2 \\
& - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\
& + \alpha \cdot \frac{1}{2\theta} \mathbb{E} \left[ \|\mathbf{y}^k - \mathbf{y}^{k-1}\|^2 \right] + \alpha \cdot \frac{1}{8\eta} \mathbb{E} \left[ \|\mathbf{z}^k - \mathbf{z}^{k-1}\|^2 \right] \\
& + \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \cdot \frac{\gamma + \frac{1}{2}\eta\mu}{p\eta} \mathbb{E} \left[ \|\mathbf{w}^k - \mathbf{z}^*\|^2 \right] + \alpha \cdot \frac{\gamma}{2\eta} \mathbb{E} \left[ \|\mathbf{x}^k - \mathbf{w}^{k-1}\|^2 \right].
\end{aligned}$$

Definition (29) of the Lyapunov function and the choice  $\alpha = \max \left[ \left( 1 - \frac{\mu\eta}{4} \right); \left( 1 - \beta\theta\chi^{-1} \right); \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \right]$  move us to

$$\mathbb{E}\Psi^{k+1} \leq \max \left[ \left( 1 - \frac{\mu\eta}{4} \right); \left( 1 - \beta\theta\chi^{-1} \right); \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \right] \mathbb{E}\Psi^k.$$

It remains to show, that

$$\Psi^k \geq \frac{1}{2\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2.$$

$$\begin{aligned} \Psi^k &\geq \frac{1}{\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{1}{8\eta} \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 + \frac{1}{2\theta} \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \\ &\quad + 2 \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^* - \mathbf{z}^k \rangle \\ &\geq \frac{1}{\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{1}{8\eta} \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 + \frac{1}{2\theta} \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \\ &\quad - \frac{1}{2\theta} \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 - 2\theta \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 - \frac{1}{8\eta L^2} \left\| \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) \right\|^2 - 8\eta L^2 \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2. \end{aligned}$$

Using  $L$ -Lipschitzness of  $\mathbf{F}$  we get

$$\Psi^k \geq \frac{1}{2\eta} (2 - 8\eta^2 L^2 - 4\theta\eta) \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2.$$

With  $\eta \leq \frac{1}{16L}$  and  $\theta \leq \frac{1}{16\eta}$  we get

$$\Psi^k \geq \frac{1}{2\eta} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2.$$

□

**Theorem D.3** (Theorem 4.1). *Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a fixed connected graph  $\mathcal{G}$  with a gossip matrix  $\mathbf{W}$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 1 with parameters*

$$\begin{aligned} \gamma = p \leq \frac{1}{8}, \quad \eta = \min \left\{ \frac{\sqrt{\gamma b}}{4L}, \frac{1}{16L\sqrt{\chi}} \right\}, \quad \beta = \min \left\{ \frac{\mu}{4L^2}; \frac{b\gamma}{4\eta\bar{L}^2} \right\}, \quad \theta = \min \left\{ \frac{1}{2\beta}; \frac{1}{16\eta} \right\}, \\ \alpha = \max \left[ \left( 1 - \frac{\mu\eta}{4} \right); \left( 1 - \beta\theta\chi^{-1} \right); \left( 1 - \frac{p\eta\mu}{2\gamma + \eta\mu} \right) \right]. \end{aligned}$$

Then, given  $\varepsilon > 0$ , the number of iterations for  $\mathbb{E}[\|\mathbf{z}^k - \mathbf{z}^*\|^2] \leq \varepsilon$  is

$$O \left( \left[ \frac{1}{p} + \chi + \frac{1}{\sqrt{pb}} \frac{\bar{L}}{\mu} + \sqrt{\chi} \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right).$$

*Proof.* It is easy to check that  $\alpha \geq \frac{1}{2}$ , then also one can verify that the choice of  $\gamma, \eta, \beta, \theta, \alpha$  satisfies the conditions of Lemma D.2. We can get that the iteration complexity of Algorithm 1:

$$\begin{aligned} O \left( \left[ 1 + \frac{1}{\eta\mu} + \frac{1}{\beta\theta\chi^{-1}} + \frac{\gamma + \eta\mu}{p\eta\mu} \right] \log \frac{1}{\varepsilon} \right) &= O \left( \left[ \frac{1}{p} + \frac{1}{\eta\mu} + \frac{1}{\beta\theta\chi^{-1}} \right] \log \frac{1}{\varepsilon} \right) \\ &= O \left( \left[ \frac{1}{p} + \chi + \frac{1}{\eta\mu} + \frac{\chi\eta}{\beta} \right] \log \frac{1}{\varepsilon} \right) \\ &= O \left( \left[ \frac{1}{p} + \chi + \frac{1}{\eta\mu} + \frac{\chi\eta^2\bar{L}^2}{bp} + \frac{\chi\eta L^2}{\mu} \right] \log \frac{1}{\varepsilon} \right) \\ &= O \left( \left[ \frac{1}{p} + \chi + \frac{1}{\sqrt{pb}} \frac{\bar{L}}{\mu} + \sqrt{\chi} \frac{L}{\mu} \right] \log \frac{1}{\varepsilon} \right). \end{aligned}$$

□

## E Proof of Theorem 4.3

We start the proof from the following lemma on  $\delta^k$  and  $\delta^{k+1/2}$  from Algorithm 2.

**Lemma E.1.** *The following inequality holds:*

$$\mathbb{E}_k \left[ \left\| \delta^k - \mathbb{E}_k [\delta^k] \right\|^2 \right] \leq \frac{2\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 + \alpha^2 \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right]. \quad (32)$$

$$\mathbb{E}_k \left[ \left\| \delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \right] \leq \frac{2\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + 2L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right]. \quad (33)$$

where  $\mathbb{E}_k [\delta^k]$  is equal to

$$\mathbb{E}_k [\delta^k] = F(\mathbf{z}^k) + \alpha(F(\mathbf{z}^k) - F(\mathbf{z}^{k-1})). \quad (34)$$

*Proof.* The proof is the same as the proof for Lemma D.1.  $\square$

Before proving the main lemma of this section, let us introduce an auxiliary notation. Let  $\hat{\mathbf{z}}^k$  be defined for all  $k = 0, 1, 2, \dots$  as follows:

$$\hat{\mathbf{x}}^k = \mathbf{x}^k - \mathbf{P}m^k, \quad (35)$$

and the Lyapunov function  $\Psi^k$  be denoted by

$$\begin{aligned} \Psi^k &= \frac{1}{\eta_y} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 + \frac{2}{\eta_x} \left\| m^k \right\|_{\mathbf{P}}^2 + \frac{1}{2\eta_y} \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \\ &+ \frac{\nu^{-1}}{\tau} \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 - 2 \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \\ &+ \frac{1}{4\eta_z} \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 + \left( 1 + \frac{3\mu\eta_z}{2} \right) \cdot \frac{1}{\eta_z} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{\omega}{2\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right]. \end{aligned} \quad (36)$$

Here we also use the next notation

$$\mathbf{y}^* = \mathbf{P}\mathbf{F}(\mathbf{z}^*) - \nu\mathbf{z}^*, \quad (37)$$

and:

$$\mathbf{x}^* = -\mathbf{P}\mathbf{F}(\mathbf{z}^*). \quad (38)$$

**Lemma E.2.** *Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a sequence of time-varying graphs  $\mathcal{G}(k)$  with gossip matrices  $\mathbf{W}(k)$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 2 with*

$$T \geq B, \quad \omega = p \leq \frac{1}{16}; \quad \theta = \frac{1}{2}; \quad \beta = 5\gamma; \quad \nu \leq \frac{\mu}{4}; \quad \tau \in (0, 1);$$

$$\eta_z \leq \min \left\{ \frac{1}{8L}, \frac{1}{32\eta_y}, \frac{\sqrt{\alpha b \omega}}{8\bar{L}} \right\}; \quad \eta_y \leq \min \left\{ \frac{1}{4\gamma}, \frac{\nu}{8\tau} \right\}; \quad \eta_x \leq \min \left\{ \frac{1}{900\chi(T)\gamma}, \frac{\nu}{36\tau\chi^2(T)} \right\};$$

$$\gamma \leq \min \left\{ \frac{\mu}{16L^2}; \frac{b\omega}{24\bar{L}^2\eta_z} \right\},$$

$$\alpha = \max \left[ \left( 1 - \frac{p\eta_z\nu}{p + \eta_z\nu} \right); 1 - \eta_y\gamma; 1 - \eta_x\gamma; 1 - \frac{\mu\eta_z}{8}; 1 - \tau; 1 - \frac{1}{4\chi(T)} \right].$$

*Let the choice of  $T$  guarantees contraction property (Assumption 2.4 point 4) with  $\chi(T)$ . Then, after  $k$  iterations we get*

$$\begin{aligned} &\mathbb{E} \left[ \frac{1}{4\eta_z} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\ &\leq \max \left[ \left( 1 - \frac{p\eta_z\nu}{\omega + \eta_z\nu} \right); 1 - \eta_y\gamma; 1 - \eta_x\gamma, \left( 1 - \frac{\mu\eta_z}{8} \right); 1 - \tau; \left( 1 - \frac{1}{4\chi(T)} \right) \right]^k \cdot \Psi^0. \end{aligned}$$

*Proof. Part 1.* We start from

$$\begin{aligned} \frac{1}{\eta_z} \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 &= \frac{1}{\eta_z} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{2}{\eta_z} \langle \mathbf{z}^{k+1} - \mathbf{z}^k, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle - \frac{1}{\eta_z} \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \\ &= \frac{1}{\eta_z} \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + \frac{2\omega}{\eta_z} \langle \mathbf{w}^k - \mathbf{z}^k, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \\ &\quad - 2 \langle \Delta_z^k - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu\mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle - \frac{1}{\eta_z} \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \end{aligned}$$

$$-2\langle \frac{1}{\eta_z}(\mathbf{z}^k + \omega(\mathbf{w}^k - \mathbf{z}^k) - \eta_z \Delta_z^k - \mathbf{z}^{k+1}) - (-\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle.$$

Optimality condition for (1)+(5) it follows, that

$$-F(z^*) \in \partial g(z^*).$$

Let us define  $\Delta^* \in (\mathbb{R}^d)^M$  as

$$\Delta^* = \frac{1}{M}[F(z^*), \dots, F(z^*)]^T.$$

It is to note that  $-\Delta^* \in M\partial \mathbf{g}(\mathbf{z}^*)$ . On the other hand with notation (37), we get

$$\Delta^* = (\mathbf{1}_M \mathbf{1}_M^\top \otimes \mathbf{I}_d) \mathbf{F}(\mathbf{z}^*) = M(\mathbf{F}(\mathbf{z}^*) - \mathbf{P}\mathbf{F}(\mathbf{z}^*)) = M(\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*)$$

It means that  $-\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^* \in \partial \mathbf{g}(\mathbf{z}^*)$ . From update for  $\mathbf{z}^{k+1}$  of Algorithm 2 it follows, that  $\frac{1}{\eta_z}(\mathbf{z}^k + \omega(\mathbf{w}^k - \mathbf{z}^k) - \mathbf{z}^{k+1} - \eta_z \Delta_z^k) \in \partial \mathbf{g}(\mathbf{z}^{k+1})$ . This together with  $-\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^* \in \partial \mathbf{g}(\mathbf{z}^*)$  and monotonicity of  $\partial \mathbf{g}(\cdot)$  implies

$$\begin{aligned} \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] &\leq \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{2\omega}{\eta_z} \mathbb{E} \left[ \langle \mathbf{w}^k - \mathbf{z}^k, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - 2\mathbb{E} \left[ \langle \Delta_z^k - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] - \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\ &= \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{2\omega}{\eta_z} \mathbb{E} \left[ \langle \mathbf{w}^k - \mathbf{z}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - \frac{2\omega}{\eta_z} \mathbb{E} \left[ \langle \mathbf{z}^k - \mathbf{z}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] - 2\mathbb{E} \left[ \langle \Delta_z^k - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\ &= \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 + \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 - \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] \\ &\quad - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 + \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 - \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\ &\quad - 2\mathbb{E} \left[ \langle \Delta_z^k - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] - \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\ &= \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] \\ &\quad - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - 2\mathbb{E} \left[ \langle \Delta_z^k - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad - \frac{1-\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right]. \end{aligned}$$

In previous we also use the simple fact  $\|a+b\|^2 = \|a\|^2 + 2\langle a, b \rangle + \|b\|^2$  twice. Definition  $\Delta_z^k$  and small rearrangement gives

$$\begin{aligned} \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] &\leq \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] \\ &\quad - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - \frac{1-\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\ &\quad - 2\mathbb{E} \left[ \langle \mathbb{E} \left[ \delta^k \right] - \nu \mathbf{z}^k - \mathbf{y}^k - \alpha(\mathbf{y}^k - \mathbf{y}^{k-1}) - (\mathbf{F}(\mathbf{z}^*) - \mathbf{y}^* - \nu \mathbf{z}^*), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ &\quad + 2\mathbb{E} \left[ \langle \mathbb{E} \left[ \delta^k \right] - \delta^k, \mathbf{z}^{k+1} - \mathbf{z}^k \rangle \right] + 2\mathbb{E} \left[ \langle \mathbb{E} \left[ \delta^k \right] - \delta^k, \mathbf{z}^k - \mathbf{z}^* \rangle \right]. \end{aligned}$$

Using the tower property of expectation we can obtain the following:

$$\begin{aligned} \mathbb{E} \left[ \langle \mathbb{E}_k \left[ \delta^k \right] - \delta^k, \mathbf{z}^k - \mathbf{z}^* \rangle \right] &= \mathbb{E} \left[ \mathbb{E}_k \left[ \langle \mathbb{E}_k \left[ \delta^k \right] - \delta^k, \mathbf{z}^k - \mathbf{z}^* \rangle \right] \right] \\ &= \mathbb{E} \left[ \langle \mathbb{E}_k \left[ \mathbb{E}_k \left[ \delta^k \right] - \delta^k \right], \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ &= \mathbb{E} \left[ \langle \mathbb{E}_k \left[ \delta^k \right] - \mathbb{E}_k \left[ \delta^k \right], \mathbf{z}^k - \mathbf{z}^* \rangle \right] = 0. \end{aligned}$$

Then we have the following

$$\frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \leq \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right]$$









$$+ \frac{\alpha}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] + \frac{4\eta_z \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right].$$

Choice of  $\omega \leq \frac{1}{16}$  and small rearrangement give

$$\begin{aligned} & \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (2\mu - \nu) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] \\ & - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^k) - (\mathbf{y}^{k+1} - \mathbf{y}^k), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ & \leq \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - \left( \frac{\omega}{\eta_z} - \nu \right) \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\ & + \alpha \cdot \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ & + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + 2\mathbb{E} \left[ \langle \mathbf{y}^{k+1} - \mathbf{y}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ & + \frac{\alpha}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] + \frac{4\eta_z \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right]. \end{aligned}$$

**Part 2.** Update for  $m^{k+1}$  and Assumption 2.4 on time-varying graph give

$$\begin{aligned} \left\| m^{k+1} \right\|_{\mathbf{P}}^2 & \leq (1 - \chi^{-1}(T)) \left\| m^k + \eta_x \Delta_x^k \right\|_{\mathbf{P}}^2 \\ & \leq (1 - \chi^{-1}(T)) \left( (1 + (2\chi(T))^{-1}) \left\| m^k \right\|_{\mathbf{P}}^2 + (1 + 2\chi(T)) \left\| \eta_x \Delta_x^k \right\|_{\mathbf{P}}^2 \right) \\ & \leq (1 - (2\chi(T))^{-1}) \left\| m^k \right\|_{\mathbf{P}}^2 + 2\eta_x^2 \chi(T) \left\| \Delta_x^k \right\|_{\mathbf{P}}^2. \end{aligned}$$

After rearranging we get

$$\left\| m^k \right\|_{\mathbf{P}}^2 \leq (1 - (4\chi(T))^{-1}) 4\chi(T) \left\| m^k \right\|_{\mathbf{P}}^2 - 4\chi(T) \left\| m^{k+1} \right\|_{\mathbf{P}}^2 + 8\eta_x^2 \chi^2(T) \left\| \Delta_x^k \right\|_{\mathbf{P}}^2. \quad (39)$$

**Part 3.** Updates for  $\mathbf{x}^{k+1}$  and  $m^{k+1}$  of Algorithm 2 imply  $\hat{\mathbf{x}}^{k+1} = \hat{\mathbf{x}}^k - \eta_x \mathbf{P} \Delta_x^k$ . By this together with update for  $\mathbf{y}^{k+1}$  of Algorithm 2 we obtain

$$\begin{aligned} & \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \\ & = \frac{1}{\eta_y} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \hat{\mathbf{x}}^k \right\|^2 \\ & + \frac{2}{\eta_y} \langle \mathbf{y}^{k+1} - \mathbf{y}^k, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle + \frac{2}{\eta_x} \langle \hat{\mathbf{x}}^{k+1} - \hat{\mathbf{x}}^k, \hat{\mathbf{x}}^k - \mathbf{x}^* \rangle \\ & \leq \frac{1}{\eta_y} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\ & - 2\langle \Delta_y^k, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - 2\langle \mathbf{P} \Delta_x^k, \mathbf{x}^k - \mathbf{x}^* \rangle + 2 \left\| \Delta_x^k \right\|_{\mathbf{P}} \left\| m^k \right\|_{\mathbf{P}} + \eta_x \left\| \Delta_x^k \right\|_{\mathbf{P}}^2. \end{aligned}$$

Using the definitions of  $\Delta_y^k$  and  $\Delta_x^k$  we get

$$\begin{aligned} & \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \\ & \leq \frac{1}{\eta_y} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 + 2 \left\| \Delta_x^k \right\|_{\mathbf{P}} \left\| m^k \right\|_{\mathbf{P}} + \eta_x \left\| \Delta_x^k \right\|_{\mathbf{P}}^2 \\ & - 2\langle \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \mathbf{z}^{k+1} + \gamma(\mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\ & - 2\langle \nu^{-1} \mathbf{P}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \beta \mathbf{P}(\mathbf{x}^k + \Delta^{k+1/2}), \mathbf{x}^k - \mathbf{x}^* \rangle \\ & = \frac{1}{\eta_y} \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 + 2 \left\| \Delta_x^k \right\|_{\mathbf{P}} \left\| m^k \right\|_{\mathbf{P}} + \eta_x \left\| \Delta_x^k \right\|_{\mathbf{P}}^2 \\ & - 2\langle \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \mathbf{z}^{k+1} + \gamma(\mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\ & - 2\langle \nu^{-1} \mathbf{P}(\mathbf{y}_c^k + \mathbf{x}_c^k), \mathbf{x}^k - \mathbf{x}^* \rangle - 2\langle \beta \mathbf{P}(\mathbf{x}^k + \Delta^{k+1/2}), \mathbf{x}^k - \mathbf{x}^* \rangle. \end{aligned}$$

Definitions (37), (38) gives  $\mathbf{P}\mathbf{F}(\mathbf{z}^*) + \mathbf{x}^* = 0$ ,  $\mathbf{P}(\mathbf{y}^* + \mathbf{x}^*) = 0$ ,  $\nu^{-1}(\mathbf{y}^* + \mathbf{x}^*) + \mathbf{z}^* = 0$ . Additionally, using  $\mathbf{P}\mathbf{x}^k = \mathbf{x}^k$ , we get

$$\frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2$$

$$\begin{aligned}
&\leq \frac{1}{\eta_y} \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
&\quad - 2\langle \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \mathbf{z}^{k+1} + \gamma(\mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - 2\langle \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k), \mathbf{x}^k - \mathbf{x}^* \rangle - 2\langle \beta(\mathbf{x}^k + \mathbf{P}\Delta^{k+1/2}), \mathbf{x}^k - \mathbf{x}^* \rangle \\
&= \frac{1}{\eta_y} \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
&\quad - 2\langle \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + \mathbf{z}^{k+1} - \nu^{-1}(\mathbf{y}^* + \mathbf{x}^*) - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - 2\gamma\langle \mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k - (\mathbf{y}^* + \mathbf{x}^* + \nu \mathbf{z}^*), \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - 2\langle \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*), \mathbf{x}^k - \mathbf{x}^* \rangle \\
&\quad - 2\beta\langle \mathbf{x}^k + \mathbf{P}\Delta^{k+1/2} - \mathbf{P}\mathbf{F}(\mathbf{z}^*) - \mathbf{x}^*, \mathbf{x}^k - \mathbf{x}^* \rangle.
\end{aligned}$$

And then

$$\begin{aligned}
&\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
&\leq \frac{1}{\eta_y} \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
&\quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - 2\nu^{-1}\langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^{k+1} + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle \\
&\quad - 2\gamma\langle \mathbf{y}^k + \mathbf{x}^k + \nu \mathbf{z}^k - \mathbf{y}^* - \mathbf{x}^* - \nu \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - 2\beta\langle \mathbf{x}^k + \mathbf{P}\Delta^{k+1/2} - \mathbf{x}^* - \mathbf{P}\mathbf{F}(\mathbf{z}^*), \mathbf{x}^k - \mathbf{x}^* \rangle.
\end{aligned}$$

Using parallelogram rule we get

$$\begin{aligned}
&\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
&\leq \frac{1}{\eta_y} \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
&\quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - 2\nu^{-1}\langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^{k+1} + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle \\
&\quad - \gamma \left( \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 - \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \right) \\
&\quad + \gamma \left( \|\mathbf{x}^k - \mathbf{x}^* + \nu(\mathbf{z}^k - \mathbf{z}^*)\|^2 + \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 \right) \\
&\quad - 2\beta \|\mathbf{x}^k - \mathbf{x}^*\|^2 + \beta \|\mathbf{x}^k - \mathbf{x}^*\|^2 + \beta \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 \\
&\leq \frac{1}{\eta_y} \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left( \frac{1}{\eta_y} - \gamma \right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
&\quad + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
&\quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - 2\nu^{-1}\langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^{k+1} + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle \\
&\quad - \gamma \|\mathbf{y}^k - \mathbf{y}^*\|^2 - (\beta - 2\gamma) \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + \beta \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2.
\end{aligned}$$

The definition of  $\beta = 5\gamma$  gives

$$\begin{aligned}
&\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
&\leq \frac{1}{\eta_y} \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left( \frac{1}{\eta_y} - \gamma \right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
&\quad + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
&\quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - 2\nu^{-1}\langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^{k+1} + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle \\
&\quad - \gamma \|\mathbf{y}^k - \mathbf{y}^*\|^2 - 3\gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2.
\end{aligned}$$

Using the definition (35) of  $\hat{\mathbf{x}}^k$ , we get

$$\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2$$



$$\begin{aligned}
&= \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left(\frac{1}{\eta_x} - \gamma\right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
&\quad + 2\gamma \|m^k\|_{\mathbf{P}}^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 \\
&\quad - \frac{2\theta\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{(\mathbf{W}^{(k)} \otimes \mathbf{I}_d)}^2 \\
&\quad - \frac{2\nu^{-1}}{\tau} \langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}_c^k - \mathbf{x}_c^k \rangle \\
&\quad - 2\nu^{-1} \langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^k + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle
\end{aligned}$$

By parallelogram rule we obtain

$$\begin{aligned}
&\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left(\frac{1}{\eta_x} - \gamma\right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
&\quad + 2\gamma \|m^k\|_{\mathbf{P}}^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 \\
&\quad - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 + \frac{\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
&\quad + \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}_c^k - \mathbf{x}_c^k\|^2 \\
&\quad - \frac{2\theta\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{(\mathbf{W}^{(k)} \otimes \mathbf{I}_d)}^2 - 2\nu^{-1} \langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^k + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle \\
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left(\frac{1}{\eta_x} - \gamma\right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
&\quad + 2\gamma \|m^k\|_{\mathbf{P}}^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 \\
&\quad - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 + \frac{\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
&\quad + \frac{2\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} - \mathbf{y}_c^k\|^2 + \frac{2\nu^{-1}}{\tau} \|\mathbf{x}_f^{k+1} - \mathbf{x}_c^k\|^2 - \frac{2\theta\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{(\mathbf{W}^{(k)} \otimes \mathbf{I}_d)}^2 \\
&\quad - 2\nu^{-1} \langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^k + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle
\end{aligned}$$

Again by expressions for  $\mathbf{x}_f^{k+1}$  and  $\mathbf{y}_f^{k+1}$

$$\begin{aligned}
&\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left(\frac{1}{\eta_x} - \gamma\right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left(\frac{1}{\eta_y} - \gamma\right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
&\quad + 2\gamma \|m^k\|_{\mathbf{P}}^2 + 2 \|\Delta_x^k\|_{\mathbf{P}} \|m^k\|_{\mathbf{P}} + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{x}^*)\|^2 \\
&\quad - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 + \frac{\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
&\quad + 2\nu^{-1} \tau \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 + \frac{2\theta^2\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{(\mathbf{W}^{(k)} \otimes \mathbf{I}_d)}^2 - \frac{2\theta\nu^{-1}}{\tau} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{(\mathbf{W}^{(k)} \otimes \mathbf{I}_d)}^2 \\
&\quad - 2\nu^{-1} \langle \mathbf{y}_c^k + \mathbf{x}_c^k - \mathbf{y}^* - \mathbf{x}^*, \mathbf{y}^k + \mathbf{x}^k - \mathbf{y}^* - \mathbf{x}^* \rangle.
\end{aligned}$$

Using the contraction property of the gossip matrix and the definition of  $\theta = \frac{1}{2}$  we get

$$\frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2$$



$$\begin{aligned}
& -2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 \\
& - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 + \frac{\nu^{-1}(1-\tau)}{\tau} \|\mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^*\|^2.
\end{aligned}$$

Using Young's inequality we get

$$\begin{aligned}
& \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
& \quad + 2\gamma \|m^k\|_{\mathbf{P}}^2 + 4\eta_x\chi(T) \|\Delta_x^k\|_{\mathbf{P}}^2 + (4\eta_x\chi(T))^{-1} \|m^k\|_{\mathbf{P}}^2 + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
& \quad - \frac{\nu^{-1}}{2\tau\chi(T)} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 \\
& \quad + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
& \quad + \frac{\nu^{-1}(1-\tau)}{\tau} \|\mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
& = \left( \frac{1}{\eta_y} - \gamma \right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
& \quad + 2\gamma \|m^k\|_{\mathbf{P}}^2 + 4\eta_x\chi(T) \|\Delta_x^k\|_{\mathbf{P}}^2 + (4\eta_x\chi(T))^{-1} \|m^k\|_{\mathbf{P}}^2 + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
& \quad - \frac{\nu^{-1}}{2\tau\chi(T)} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 \\
& \quad + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
& \quad + \frac{\nu^{-1}(1-\tau)}{\tau} \|\mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^*\|^2.
\end{aligned}$$

By the assumption on  $\eta_x \leq \frac{1}{8\chi(T)\gamma}$  we get that  $2\gamma \leq \frac{1}{4\eta_x}$  and

$$\begin{aligned}
& \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
& \quad + 4\eta_x\chi(T) \|\Delta_x^k\|_{\mathbf{P}}^2 + (2\eta_x\chi(T))^{-1} \|m^k\|_{\mathbf{P}}^2 + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 - \frac{\nu^{-1}}{2\tau\chi(T)} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{\mathbf{P}}^2 \\
& \quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 \\
& \quad + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
& \quad + \frac{\nu^{-1}(1-\tau)}{\tau} \|\mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^*\|^2.
\end{aligned}$$

Using (39) we get

$$\begin{aligned}
& \frac{1}{\eta_y} \|\mathbf{y}^{k+1} - \mathbf{y}^*\|^2 + \frac{1}{\eta_x} \|\hat{\mathbf{x}}^{k+1} - \mathbf{x}^*\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \|\mathbf{y}^k - \mathbf{y}^*\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \|\hat{\mathbf{x}}^k - \mathbf{x}^*\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \|\mathbf{y}^{k+1} - \mathbf{y}^k\|^2 \\
& \quad + 8\eta_x\chi(T) \|\Delta_x^k\|_{\mathbf{P}}^2 + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \|m^k\|_{\mathbf{P}}^2 - \frac{2}{\eta_x} \|m^{k+1}\|_{\mathbf{P}}^2 + \eta_x \|\Delta_x^k\|_{\mathbf{P}}^2 \\
& \quad - \frac{\nu^{-1}}{2\tau\chi(T)} \|\mathbf{y}_c^k + \mathbf{x}_c^k\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \|\mathbf{x}^k - \mathbf{x}^*\|^2 + 2\gamma\nu^2 \|\mathbf{z}^k - \mathbf{z}^*\|^2 \\
& \quad + 5\gamma \|\Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*)\|^2 - \frac{\nu^{-1}}{\tau} \|\mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^*\|^2 \\
& \quad + \frac{\nu^{-1}(1-\tau)}{\tau} \|\mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^*\|^2.
\end{aligned}$$

Using the definition of  $\Delta_x^k$  with definition of  $\beta = 5\gamma$  we get

$$\begin{aligned}
& \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
& \quad + 9\eta_x \chi(T) \left\| \nu^{-1}(\mathbf{y}_c^k + \mathbf{x}_c^k) + 5\gamma(\mathbf{x}^k + \Delta^{k+1/2}) \right\|_{\mathbf{P}}^2 + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \left\| m^k \right\|_{\mathbf{P}}^2 \\
& \quad - \frac{2}{\eta_x} \left\| m^{k+1} \right\|_{\mathbf{P}}^2 - \frac{\nu^{-1}}{2\tau\chi(T)} \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \left\| \mathbf{x}^k - \mathbf{x}^* \right\|^2 \\
& \quad + 2\gamma\nu^2 \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + 5\gamma \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 - \frac{\nu^{-1}}{\tau} \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \\
& \quad + \frac{\nu^{-1}(1-\tau)}{\tau} \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2.
\end{aligned}$$

By definition  $\mathbf{P}\mathbf{F}(\mathbf{z}^*) + \mathbf{x}^* = 0$  we obtain

$$\begin{aligned}
& \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
& \quad + 18\eta_x \nu^{-2} \chi(T) \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 + 450\eta_x \chi(T) \gamma^2 \left\| \mathbf{x}^k - \mathbf{x}^* - \mathbf{P}\mathbf{F}(\mathbf{z}^*) + \Delta^{k+1/2} \right\|_{\mathbf{P}}^2 \\
& \quad + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \left\| m^k \right\|_{\mathbf{P}}^2 - \frac{2}{\eta_x} \left\| m^{k+1} \right\|_{\mathbf{P}}^2 - \frac{\nu^{-1}}{2\tau\chi(T)} \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 \\
& \quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \left\| \mathbf{x}^k - \mathbf{x}^* \right\|^2 + 2\gamma\nu^2 \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + 5\gamma \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \\
& \quad - \frac{\nu^{-1}}{\tau} \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 + \frac{\nu^{-1}(1-\tau)}{\tau} \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
& \quad + 18\eta_x \nu^{-2} \chi(T) \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 + 900\eta_x \chi(T) \gamma^2 \left\| \mathbf{x}^k - \mathbf{x}^* \right\|_{\mathbf{P}}^2 \\
& \quad + 900\eta_x \chi(T) \gamma^2 \left\| \Delta^{k+1/2} - \mathbf{P}\mathbf{F}(\mathbf{z}^*) \right\|_{\mathbf{P}}^2 + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \left\| m^k \right\|_{\mathbf{P}}^2 \\
& \quad - \frac{2}{\eta_x} \left\| m^{k+1} \right\|_{\mathbf{P}}^2 - \frac{\nu^{-1}}{2\tau\chi(T)} \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \left\| \mathbf{x}^k - \mathbf{x}^* \right\|^2 \\
& \quad + 2\gamma\nu^2 \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + 5\gamma \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \\
& \quad - \frac{\nu^{-1}}{\tau} \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 + \frac{\nu^{-1}(1-\tau)}{\tau} \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2.
\end{aligned}$$

With the assumption on  $\eta_x \leq \frac{1}{900\chi(T)\gamma}; \frac{\nu}{36\tau\chi^2(T)}$  we get

$$\begin{aligned}
& \frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \\
& \leq \left( \frac{1}{\eta_y} - \gamma \right) \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \left( \frac{1}{\eta_x} - \gamma \right) \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \left( \frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau \right) \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
& \quad + \frac{\nu^{-1}}{2\tau\chi(T)} \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 + \gamma \left\| \mathbf{x}^k - \mathbf{x}^* \right\|_{\mathbf{P}}^2 + \gamma \left\| \Delta^{k+1/2} - \mathbf{P}\mathbf{F}(\mathbf{z}^*) \right\|_{\mathbf{P}}^2 \\
& \quad + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \left\| m^k \right\|_{\mathbf{P}}^2 - \frac{2}{\eta_x} \left\| m^{k+1} \right\|_{\mathbf{P}}^2 - \frac{\nu^{-1}}{2\tau\chi(T)} \left\| \mathbf{y}_c^k + \mathbf{x}_c^k \right\|_{\mathbf{P}}^2 \\
& \quad - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle - \gamma \left\| \mathbf{x}^k - \mathbf{x}^* \right\|^2 + 2\gamma\nu^2 \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + 5\gamma \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \\
& \quad - \frac{\nu^{-1}}{\tau} \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 + \frac{\nu^{-1}(1-\tau)}{\tau} \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2.
\end{aligned}$$

With property of the projector:  $\mathbf{P}\mathbf{P} = \mathbf{P}$ , we get

$$\frac{1}{\eta_y} \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 + \frac{1}{\eta_x} \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2$$

$$\begin{aligned}
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 + \left(\frac{1}{\eta_x} - \gamma\right) \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 - \left(\frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau\right) \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \\
&\quad + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \left\| m^k \right\|_{\mathbf{P}}^2 - \frac{2}{\eta_x} \left\| m^{k+1} \right\|_{\mathbf{P}}^2 - 2\langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \\
&\quad + 2\gamma\nu^2 \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 + 6\gamma \left\| \Delta^{k+1/2} - \mathbf{F}(\mathbf{z}^*) \right\|^2 \\
&\quad - \frac{\nu^{-1}}{\tau} \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 + \frac{\nu^{-1}(1-\tau)}{\tau} \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2.
\end{aligned}$$

Taking full expectation and using (33) we have

$$\begin{aligned}
&\frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] \\
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + \left(\frac{1}{\eta_x} - \gamma\right) \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] \\
&\quad - \left(\frac{1}{\eta_y} - \gamma - 2\nu^{-1}\tau\right) \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] \\
&\quad - \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \right] + 2\gamma\nu^2 \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
&\quad + 12\gamma L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{12\gamma \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] \\
&\quad - \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] + \frac{\nu^{-1}(1-\tau)}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right].
\end{aligned}$$

Using the definition of  $\eta_y$  we get that  $\gamma \leq \frac{1}{4\eta_y}$  and  $2\nu^{-1}\tau \leq \frac{1}{4\eta_y}$

$$\begin{aligned}
&\frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] \\
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + \left(\frac{1}{\eta_x} - \gamma\right) \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] - \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\
&\quad + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] - \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \right] \\
&\quad + 2\gamma\nu^2 \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + 12\gamma L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{12\gamma \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] \\
&\quad - \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] + \frac{\nu^{-1}(1-\tau)}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right].
\end{aligned}$$

**Part 4.** After combining parts 1 and 3 of this proof we get

$$\begin{aligned}
&\frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] + \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + (2\mu - \nu) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\
&\quad + \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^k) - (\mathbf{y}^{k+1} - \mathbf{y}^k), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\
&\leq \left(\frac{1}{\eta_y} - \gamma\right) \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + \left(\frac{1}{\eta_x} - \gamma\right) \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] - \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\
&\quad + (1 - (4\chi(T))^{-1}) \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] - \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{z}^{k+1} - \mathbf{z}^*, \mathbf{y}^{k+1} - \mathbf{y}^* \rangle \right] \\
&\quad + 2\gamma\nu^2 \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + 12\gamma L^2 \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] + \frac{12\gamma \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] \\
&\quad - \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] + \frac{\nu^{-1}(1-\tau)}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\
&\quad + \frac{1}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] - \left(\frac{\omega}{\eta_z} - \nu\right) \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\
&\quad + \alpha \cdot \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\
&\quad + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right] - \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + 2\mathbb{E} \left[ \langle \mathbf{y}^{k+1} - \mathbf{y}^*, \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right]
\end{aligned}$$



$$+ \frac{\alpha}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] + \frac{4\eta_z \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right].$$

Small rearrangement gives

$$\begin{aligned} & \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] + \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\ & + \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^k) - (\mathbf{y}^{k+1} - \mathbf{y}^k), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ & + \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] + \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \left( 1 - \frac{12\gamma \bar{L}^2 \eta_z}{b\omega} \right) \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + \left( \frac{1}{\eta_z} + 2\mu - \nu - 12\gamma L^2 \right) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & \leq (1 - \eta_y \gamma) \cdot \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + (1 - \eta_x \gamma) \cdot \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] + \alpha \cdot \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\ & + \alpha \cdot \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ & + (1 - (4\chi(T))^{-1}) \cdot \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] + (1 - \tau) \cdot \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \frac{4\eta_z \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] + \left( \frac{1}{\eta_z} - \frac{\omega}{\eta_z} + \nu + 2\gamma \nu^2 \right) \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\ & + \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right]. \end{aligned}$$

Now, we add  $\frac{\gamma + \eta \nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right]$  to both sides and use update for  $\mathbf{w}^{k+1}$

$$\frac{\omega + \eta_z \nu}{p\eta_z} \mathbb{E} \left[ \mathbb{E}_{\mathbf{w}^{k+1}} \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right] = \frac{\omega + \eta_z \nu}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] + \frac{(\omega + \eta_z \nu)(1 - p)}{\eta_z p} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right],$$

and then

$$\begin{aligned} & \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] + \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\ & + \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^k) - (\mathbf{y}^{k+1} - \mathbf{y}^k), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ & + \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] + \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \left( 1 - \frac{12\gamma \bar{L}^2 \eta_z}{b\omega} \right) \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + \left( \frac{1}{\eta_z} + 2\mu - \nu - 12\gamma L^2 \right) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & + \frac{\omega + \eta_z \nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & \leq (1 - \eta_y \gamma) \cdot \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + (1 - \eta_x \gamma) \cdot \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] + \alpha \cdot \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\ & + \alpha \cdot \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ & + (1 - (4\chi(T))^{-1}) \cdot \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] + (1 - \tau) \cdot \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \frac{4\eta_z \bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] + \left( \frac{1}{\eta_z} + 2\nu + 2\gamma \nu^2 \right) \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\ & + \left( 1 - \frac{p\eta_z \nu}{\omega + \eta_z \nu} \right) \cdot \frac{\omega + \eta_z \nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right]. \end{aligned}$$

With  $\gamma \leq \frac{\mu}{48L^2}$  and  $\nu \leq \frac{\mu}{4}$  we get that

$$(1 + 2\mu\eta_x - \nu\eta_x - 12\gamma L^2 \eta_x) \geq 1 + \frac{3\mu\eta_x}{2},$$

$$\left(\frac{1}{\eta_x} + 2\nu + 2\gamma\nu^2\right) \leq \left(1 + \frac{\mu\eta_x}{2} + \frac{2\mu^3\eta_x}{16 \cdot 48L^2}\right) \frac{1}{\eta_x} \leq (1 + \mu\eta_x) \frac{1}{\eta_x}.$$

Additionally, by

$$(1 + \mu\eta_x) \leq \left(1 + \frac{3\mu\eta_x}{2}\right) \left(1 - \frac{\mu\eta_x}{8}\right)$$

we obtain

$$\begin{aligned} & \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] + \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\ & + \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^k) - (\mathbf{y}^{k+1} - \mathbf{y}^k), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ & + \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] + \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \left(1 - \frac{12\gamma\bar{L}^2\eta_z}{b\omega}\right) \frac{\omega}{\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + \left(1 + \frac{3\mu\eta_x}{2}\right) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & + \frac{\omega + \eta_z\nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & \leq (1 - \eta_y\gamma) \cdot \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + (1 - \eta_x\gamma) \cdot \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] + \alpha \cdot \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\ & + \alpha \cdot \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ & + (1 - (4\chi(T))^{-1}) \cdot \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] + (1 - \tau) \cdot \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \frac{4\eta_z\bar{L}^2}{b} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] + \left(1 - \frac{\mu\eta_x}{8}\right) \cdot \left(1 + \frac{3\mu\eta_x}{2}\right) \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\ & + \left(1 - \frac{p\eta_z\nu}{\omega + \eta_z\nu}\right) \cdot \frac{\omega + \eta_z\nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right]. \end{aligned}$$

The choice  $\eta_z \leq \frac{\sqrt{\alpha\omega b}}{4L}$  and  $\gamma \leq \frac{b\omega}{24L^2\eta_z}$  gives

$$\begin{aligned} & \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^* \right\|^2 \right] + \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^{k+1} - \mathbf{x}^* \right\|^2 \right] + \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^{k+1} - \mathbf{y}^k \right\|^2 \right] \\ & + \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^k \right\|^2 \right] - 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^{k+1}) - \mathbf{F}(\mathbf{z}^k) - (\mathbf{y}^{k+1} - \mathbf{y}^k), \mathbf{z}^{k+1} - \mathbf{z}^* \rangle \right] \\ & + \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^{k+1} \right\|_{\mathbf{P}}^2 \right] + \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^{k+1} + \mathbf{x}_f^{k+1} - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \frac{\omega}{2\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{w}^k \right\|^2 \right] + \left(1 + \frac{3\mu\eta_x}{2}\right) \mathbb{E} \left[ \left\| \mathbf{z}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & + \frac{\omega + \eta_z\nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^{k+1} - \mathbf{z}^* \right\|^2 \right] \\ & \leq (1 - \eta_y\gamma) \cdot \frac{1}{\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^* \right\|^2 \right] + (1 - \eta_x\gamma) \cdot \frac{1}{\eta_x} \mathbb{E} \left[ \left\| \hat{\mathbf{x}}^k - \mathbf{x}^* \right\|^2 \right] + \alpha \cdot \frac{1}{2\eta_y} \mathbb{E} \left[ \left\| \mathbf{y}^k - \mathbf{y}^{k-1} \right\|^2 \right] \\ & + \alpha \cdot \frac{1}{4\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^{k-1} \right\|^2 \right] - \alpha \cdot 2\mathbb{E} \left[ \langle \mathbf{F}(\mathbf{z}^k) - \mathbf{F}(\mathbf{z}^{k-1}) - (\mathbf{y}^k - \mathbf{y}^{k-1}), \mathbf{z}^k - \mathbf{z}^* \rangle \right] \\ & + (1 - (4\chi(T))^{-1}) \cdot \frac{2}{\eta_x} \mathbb{E} \left[ \left\| m^k \right\|_{\mathbf{P}}^2 \right] + (1 - \tau) \cdot \frac{\nu^{-1}}{\tau} \mathbb{E} \left[ \left\| \mathbf{y}_f^k + \mathbf{x}_f^k - \mathbf{y}^* - \mathbf{x}^* \right\|^2 \right] \\ & + \alpha \cdot \frac{\omega}{2\eta_z} \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{w}^{k-1} \right\|^2 \right] + \left(1 - \frac{\mu\eta_x}{8}\right) \cdot \left(1 + \frac{3\mu\eta_x}{2}\right) \mathbb{E} \left[ \left\| \mathbf{z}^k - \mathbf{z}^* \right\|^2 \right] \\ & + \left(1 - \frac{p\eta_z\nu}{\omega + \eta_z\nu}\right) \cdot \frac{\omega + \eta_z\nu}{p\eta_z} \mathbb{E} \left[ \left\| \mathbf{w}^k - \mathbf{z}^* \right\|^2 \right]. \end{aligned}$$

Putting  $p = \omega$  and using definition of the Lyapunov function, we get

$$\mathbb{E}[\Psi^{k+1}] \leq \max \left[ \left(1 - \frac{p\eta_z\nu}{p + \eta_z\nu}\right); 1 - \eta_y\gamma; 1 - \eta_x\gamma; 1 - \frac{\mu\eta_z}{8}; 1 - \tau; 1 - \frac{1}{4\chi(T)} \right] \cdot \mathbb{E}[\Psi^k].$$

Finally,

$$\begin{aligned}
\Psi^k &\geq \frac{1}{\eta_x} \|x^k - \mathbf{x}^*\|^2 + \frac{1}{4\eta_x} \|x^{k-1} - x^k\|^2 + \frac{1}{2\eta_y} \|y^{k-1} - y^k\|^2 \\
&\quad - 2\langle \mathbf{F}(x^k) - \mathbf{F}(x^{k-1}) - (y^k - y^{k-1}), x^k - \mathbf{x}^* \rangle \\
&\geq \frac{1}{\eta_x} \|x^k - \mathbf{x}^*\|^2 + \frac{1}{4\eta_x} \|x^{k-1} - x^k\|^2 + \frac{1}{2\eta_y} \|y^{k-1} - y^k\|^2 \\
&\quad - \frac{1}{2\eta_y} \|y^k - y^{k-1}\|^2 - 2\eta_y \|x^k - \mathbf{x}^*\|^2 - \frac{1}{4\eta_x} \|x^{k-1} - x^k\|^2 - 4\eta_x L^2 \|x^k - \mathbf{x}^*\|^2 \\
&= (1 - 2\eta_x \eta_y - 4\eta_x^2 L^2) \frac{1}{\eta_x} \|x^k - \mathbf{x}^*\|^2 \\
&\geq \frac{1}{4\eta_x} \|x^k - \mathbf{x}^*\|^2.
\end{aligned}$$

□

**Theorem E.3** (Theorem 4.3). *Consider the problem (10) (or (1) + (5)) under Assumptions 2.1 and 2.2 over a sequence of time-varying graphs  $\mathcal{G}(k)$  with gossip matrices  $\mathbf{W}(k)$ . Let  $\{\mathbf{z}^k\}$  be the sequence generated by Algorithm 2 with parameters*

$$\begin{aligned}
T &\geq B; \quad \omega = p \leq \frac{1}{16}; \quad \theta = \frac{1}{2}; \quad \beta = 5\gamma; \quad \nu = \frac{\mu}{4}; \quad \tau = \min \left\{ \frac{\mu}{32L\chi(T)}, \frac{\mu\sqrt{bp}}{32\bar{L}} \right\}; \\
\eta_z &= \min \left\{ \frac{1}{8L\chi(T)}, \frac{1}{32\eta_y}, \frac{\sqrt{\alpha b\omega}}{8\bar{L}} \right\}; \quad \eta_y = \min \left\{ \frac{1}{4\gamma}, \frac{\nu}{8\tau} \right\}; \quad \eta_x = \min \left\{ \frac{1}{900\chi(T)\gamma}, \frac{\nu}{36\tau\chi^2(T)} \right\}; \\
\gamma &= \min \left\{ \frac{\mu}{16L^2}, \frac{b\omega}{24\bar{L}^2\eta_z} \right\}, \\
\alpha &= \max \left[ \left( 1 - \frac{p\eta_z\nu}{p + \eta_z\nu} \right); 1 - \eta_y\gamma; 1 - \eta_x\gamma; 1 - \frac{\mu\eta_z}{8}; 1 - \tau; 1 - \frac{1}{4\chi(T)} \right].
\end{aligned}$$

Let the choice of  $T$  guarantees contraction property (Assumption 2.4 point 4) with  $\chi(T)$ . Then, given  $\varepsilon > 0$ , the number of iterations for  $\mathbb{E}[\|\mathbf{z}^k - \mathbf{z}^*\|^2] \leq \varepsilon$  is

$$O \left( \left[ \chi^2(T) + \frac{1}{p} + \chi(T) \frac{L}{\mu} + \frac{1}{\sqrt{bp}} \frac{\bar{L}}{\mu} \right] \log \frac{1}{\varepsilon} \right).$$

*Proof.* It is easy to check that  $\alpha \geq \frac{1}{2}$ , then also one can verify that the choice of  $\omega, \theta, \beta, \nu, \tau, \eta_z, \eta_y, \eta_x, \gamma, \alpha$  satisfies the conditions of Lemma E.2. We can get that the iteration complexity of Algorithm 2:

$$O \left( \left[ \chi(T) + \frac{1}{p} + \frac{1}{\eta_y\gamma} + \frac{1}{\eta_x\gamma} + \frac{1}{\eta_z\mu} + \frac{1}{\eta_z\nu} + \frac{1}{\tau} + \frac{1}{1-\alpha} \right] \log \frac{1}{\varepsilon} \right).$$

Substituting  $\alpha, \nu, \omega$ , we get

$$O \left( \left[ \chi(T) + \frac{1}{p} + \frac{1}{\eta_y\gamma} + \frac{1}{\eta_x\gamma} + \frac{1}{\eta_z\mu} + \frac{1}{\tau} \right] \log \frac{1}{\varepsilon} \right).$$

Putting  $\eta_z$

$$O \left( \left[ \chi(T) + \frac{1}{p} + \chi(T) \frac{L}{\mu} + \frac{1}{\sqrt{bp}} \frac{\bar{L}}{\mu} + \frac{1}{\eta_y\gamma} + \frac{1}{\eta_x\gamma} + \frac{\eta_y}{\mu} + \frac{1}{\tau} \right] \log \frac{1}{\varepsilon} \right).$$

Substituting  $\eta_y$  and  $\eta_x$

$$O \left( \left[ \chi(T) + \frac{1}{p} + \chi(T) \frac{L}{\mu} + \frac{1}{\sqrt{bp}} \frac{\bar{L}}{\mu} + \frac{\tau\chi^2(T)}{\mu\gamma} + \frac{\eta_y}{\mu} + \frac{1}{\tau} \right] \log \frac{1}{\varepsilon} \right).$$

Using  $\eta_y \leq \frac{\nu}{8\tau}$

$$O \left( \left[ \chi(T) + \frac{1}{p} + \chi(T) \frac{L}{\mu} + \frac{1}{\sqrt{bp}} \frac{\bar{L}}{\mu} + \frac{\tau\chi^2(T)}{\mu\gamma} + \frac{1}{\tau} \right] \log \frac{1}{\varepsilon} \right).$$

Substituting  $\tau$  we get

$$O \left( \left[ \chi(T) + \frac{1}{p} + \chi(T) \frac{L}{\mu} + \frac{1}{\sqrt{bp}} \frac{\bar{L}}{\mu} + \frac{\tau\chi^2(T)}{\mu\gamma} \right] \log \frac{1}{\varepsilon} \right).$$

Putting  $\gamma$  we obtain

$$O\left(\left[\chi(T) + \frac{1}{p} + \chi(T)\frac{L}{\mu} + \frac{1}{\sqrt{bp}}\frac{\bar{L}}{\mu} + \frac{\tau\bar{L}^2\eta_z\chi^2(T)}{\mu bp} + \frac{\tau L^2\chi^2(T)}{\mu^2}\right]\log\frac{1}{\varepsilon}\right).$$

With  $\eta_z \leq \frac{\sqrt{abp}}{8L}$  and  $\tau \leq \frac{\mu\sqrt{bp}}{L}$  we have

$$O\left(\left[\chi^2(T) + \frac{1}{p} + \chi(T)\frac{L}{\mu} + \frac{1}{\sqrt{bp}}\frac{\bar{L}}{\mu} + \frac{\tau L^2\chi^2(T)}{\mu^2}\right]\log\frac{1}{\varepsilon}\right).$$

Finally,  $\tau \leq \frac{1}{8L\chi(T)}$

$$O\left(\left[\chi^2(T) + \frac{1}{p} + \chi(T)\frac{L}{\mu} + \frac{1}{\sqrt{bp}}\frac{\bar{L}}{\mu}\right]\log\frac{1}{\varepsilon}\right).$$


□

## F Additional experiments

### F.1 Variance reduction

Here we give additional experiments for Section 5.1 on other matrices.

Figure 4: Comparison epoch complexities of Algorithm 1, EG-Alc-Alg1, EG-Alc-Alg2 and EG-Car on (11) with matrices from [55] (two upper lines correspond to test matrix 1, the two bottom lines – to test matrix 2). Dashed lines give convergence with theoretical parameters, solid lines – with tuned parameters.



### F.2 Decentralized methods

#### F.2.1 Fixed networks

Here we give additional experiments for Section 5.2.1.

Figure 5: Comparison communication complexities of Algorithm 1, EGD-GT, EGD-Con and Sliding on (12) with over fixed grid networks.




Figure 6: Comparison communication complexities of Algorithm 1, EGD-GT, EGD-Con and Sliding on (12) with over fixed ring networks.





Figure 7: Comparison communication complexities of Algorithm 1, EGD-GT, EGD-Con and Sliding on (12) with over fixed star networks.



## F.2.2 Time-varying networks

Here we give additional experiments for Section 5.2.2.

Figure 8: Comparison communication complexities of Algorithm 2, EGD-GT, EGD-Con and Sliding on (12) over time-varying grid networks.





Figure 9: Comparison communication complexities of Algorithm 1, EGD-GT, EGD-Con and Sliding on (12) with over fixed ring networks.



## G Tables

Table 2: Summary communication and local complexities for finding an  $\varepsilon$ -solution for strongly monotone **deterministic decentralized** variational inequality (1) on fixed and time-varying networks. Convergence is measured by the distance to the solution. *Notation:*  $\mu$  = constant of strong monotonicity of the operator  $F$ ,  $L =$  Lipschitz constants for all  $L_{m,i}$ ,  $\chi$  = characteristic number of the network (see Assumptions 2.3 and 2.4).

		Reference	Communication complexity	Local complexity	Weaknesses
Fixed	Upper	Mukherjee and Chakraborty [48] <sup>(1,2)</sup>	$\mathcal{O}\left(\chi^{\frac{4}{3}} \frac{L}{\mu^{\frac{4}{3}}} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\chi^{\frac{4}{3}} \frac{L}{\mu^{\frac{4}{3}}} \log \frac{1}{\varepsilon}\right)$	bad comm. rates bad local rates
		Beznosikov et al. [14] <sup>(1,2)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{L+\mu}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip no linear convergence
		Beznosikov et al. [13] <sup>(1,3)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip no linear convergence
		Rogozin et al. [62] <sup>(1,2,4)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip bounded gradient
		This paper	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	
	This paper	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip	
	Lower	Beznosikov et al. [13] <sup>(3)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	
This paper	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$			
Time-varying	Upper	Beznosikov et al. [10] <sup>(3)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon} + \chi \frac{LD}{\mu^2 \sqrt{\varepsilon}}\right)$ <sup>(5)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon} + \chi \frac{LD}{\mu^2 \sqrt{\varepsilon}}\right)$	$D$ -homogeneity no linear convergence
		Beznosikov et al. [12] <sup>(1,2)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip no linear convergence
		This paper	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(5)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	
	This paper	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(5)</sup>	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	multiple gossip	
	Lower	Beznosikov et al. [12] <sup>(2)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	
This paper	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(5)</sup>	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$			

<sup>(1)</sup> for saddle point problems

<sup>(2)</sup> deterministic

<sup>(3)</sup> stochastic, but not finite sum

<sup>(4)</sup> for convex-concave (monotone) case (we reanalyzed for strongly monotone case)

<sup>(5)</sup>  $B$ -connected graphs are also considered. For simplicity in comparison with other works, we put  $B = 1$ . To get estimates for  $B \neq 1$ , one need to change  $\chi$  to  $B\chi$



Table 3: Summary complexities for finding an  $\varepsilon$ -solution for strongly monotone **stochastic (finite-sum) non-distributed** variational inequality (1). Convergence is measured by the distance to the solution. *Notation:*  $\mu$  = constant of strong monotonicity of the operator  $F$ ,  $L$  = Lipschitz constants for all  $L_i$ ,  $n$  = the size of the local dataset.

Reference	Complexity	Weaknesses
Palaniappan and Bach [59] - SVRG <sup>(1)</sup>	$\mathcal{O}\left(n + b \frac{L^2}{\mu^2} \log \frac{1}{\varepsilon}\right)$	bad complexity batching
Palaniappan and Bach [59] - Acc-SVRG <sup>(1)</sup>	$\mathcal{O}\left(n + \sqrt{bn} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	envelope acceleration batching
Chavdarova et al. [20] <sup>(2)</sup>	$\mathcal{O}\left(n + b \frac{L^2}{\mu^2} \log \frac{1}{\varepsilon}\right)$	bad complexity batching
Carmon et al. [17] <sup>(1)</sup>	$\mathcal{O}\left(n + \sqrt{bn} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	for games only batching
Carmon et al. [17] <sup>(1)</sup>	$\tilde{\mathcal{O}}\left(n + \sqrt{bn} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	batching
Yang et al. [70] <sup>(1,3)</sup>	$\mathcal{O}\left(b^{\frac{1}{3}} n^{\frac{2}{3}} \frac{L^3}{\mu^3} \log \frac{1}{\varepsilon}\right)$	bad complexity batching
Alacaoglu and Malitsky [1]	$\mathcal{O}\left(n + \sqrt{bn} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	batching
Alacaoglu et al. [2]	$\mathcal{O}\left(n + n \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	bad rates
Tominin et al. [65] <sup>(1)</sup>	$\mathcal{O}\left(n + \sqrt{bn} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	envelope acceleration batching
Beznosikov et al. [11] <sup>(2)</sup>	$\mathcal{O}\left(n + b \frac{L^2}{\mu^2} \log \frac{1}{\varepsilon}\right)$	bad complexity batching
This paper	$\mathcal{O}\left(n + \sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	

<sup>(1)</sup> for saddle point problems

<sup>(2)</sup> under  $l$ -cocoercivity assumption (in genral case  $l = \frac{L^2}{\mu}$ )

<sup>(3)</sup> under PL - condition

Table 4: Summary communication and local complexities for finding an  $\varepsilon$ -solution for strongly monotone **decentralized** variational inequality (1) on fixed and time-varying networks. Convergence is measured by the distance to the solution. *Notation:*  $\mu =$  constant of strong monotonicity of the operator  $F$ ,  $L =$  Lipschitz constants for all  $L_{m,i}$ ,  $\chi =$  characteristic number of the network (see Assumptions 2.3 and 2.4).

	Reference	Communication complexity	Local complexity	Deter/Stoch?
Fixed	Mukherjee and Chakraborty [48] <sup>(1)</sup>	$\mathcal{O}\left(\chi^{\frac{4}{3}} \frac{L^{\frac{4}{3}}}{\mu^{\frac{4}{3}}} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\chi^{\frac{4}{3}} \frac{L^{\frac{4}{3}}}{\mu^{\frac{4}{3}}} \log \frac{1}{\varepsilon}\right)$	deterministic
	Beznosikov et al. [14] <sup>(1)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{L+\mu}{\mu} \log \frac{1}{\varepsilon}\right)$	deterministic
	Beznosikov et al. [13] <sup>(1)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon} + \frac{\sigma^2}{\mu^2 M \varepsilon}\right)$	stochastic $\sigma^2$ -bounded variance
	Rogozin et al. [62] <sup>(1,2)</sup>	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	deterministic
	This paper	$\mathcal{O}\left(\max[\sqrt{n}; \sqrt{\chi}] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\max[\sqrt{n}; \sqrt{\chi}] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	stochastic finite-sum
	This paper	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	stochastic finite-sum
Lower	Beznosikov et al. [13]	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon} + \frac{\sigma^2}{\mu^2 M \varepsilon}\right)$	stochastic $\sigma^2$ -bounded variance
	This paper	$\mathcal{O}\left(\sqrt{\chi} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	stochastic finite-sum
Time-varying	Beznosikov et al. [10]	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon} + \frac{L(\chi D + \sqrt{\chi} \sigma)}{\mu^2 \sqrt{\varepsilon}} + \frac{\sigma^2}{\mu^2 M \varepsilon}\right)$ <sup>(3)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon} + \frac{L(\chi D + \sqrt{\chi} \sigma)}{\mu^2 \sqrt{\varepsilon}} + \frac{\sigma^2}{\mu^2 M \varepsilon}\right)$	$D$ -homogeneity $\sigma^2$ -bounded variance
	Beznosikov et al. [12] <sup>(1)</sup>	$\mathcal{O}\left(\chi \frac{L}{\mu} \log^2 \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	deterministic
	This paper	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(3)</sup>	$\mathcal{O}\left(\max[\sqrt{n}; \chi] \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	stochastic finite-sum
	This paper	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(3)</sup>	$\mathcal{O}\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	stochastic finite-sum
	Beznosikov et al. [12]	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	$\mathcal{O}\left(\frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	deterministic
	This paper	$\mathcal{O}\left(\chi \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$ <sup>(3)</sup>	$\mathcal{O}\left(\sqrt{n} \frac{L}{\mu} \log \frac{1}{\varepsilon}\right)$	stochastic finite-sum

<sup>(1)</sup> for saddle point problems

<sup>(2)</sup> for convex-concave (monotone) case (we reanalyzed for strongly monotone case)

<sup>(3)</sup>  $B$ -connected graphs are also considered. For simplicity in comparison with other works, we put  $B = 1$ . To get estimates for  $B \neq 1$ , one need to change  $\chi$  to  $B\chi$